arXiv:1302.6678v3 [math.AG] 12 Mar 2013

# Algorithms for singularities and real structures of weak Del Pezzo surfaces

Niels Lubbes

March 13, 2013

## Abstract

In this paper we consider the classification of singularities (Du Val) and real structures (Wall) of weak Del Pezzo surfaces from an algorithmic point of view.

It is well known that the singularities of weak Del Pezzo surfaces correspond to root subsystems. We present an algorithm which computes the classification of these root subsystems. We represent equivalence classes of root subsystems by unique labels. These labels allow us to construct examples of weak Del Pezzo surfaces with the corresponding singularity configuration.

Equivalence classes of real structures of weak Del Pezzo surfaces are also represented by root subsystems. We present an algorithm which computes the classification of real structures. This leads to an alternative proof of the known classification. As an application we classify families of real conics of weak Del Pezzo surfaces of degree four.

# Contents

1 Introduction 3

2 Root subsystems 7

3 Algorithm: geometric bases of C1 root subsystems 10

4 Weak Del Pezzo surfaces 11

5 Algorithm: analyzing and constructing linear series 16

6 Examples of constructing weak Del Pezzo surfaces 20

7 Algorithm: real structures of weak Del Pezzo surfaces 22

8 Acknowledgements 26

A Table of geometric C1 labels 27

B Real structures of weak Del Pezzo surfaces 29

# 1 Introduction

## 1.1 Problems

A weak Del Pezzo surface is an algebraic surface such that the anticanonical divisor class is nef and big. We refer to Dolgachev [2012] for the theory of weak Del Pezzo surfaces. Many properties of weak Del Pezzo surfaces are determined by its Picard group and the effective minus two classes therein. The minus two classes of a given weak Del Pezzo surface form a root system and the effective minus two classes form a root subsystem. These root subsystems are well studied, and the classification can for example be found in Du Val [1934] and Dolgachev [2012]. In Wall [1987] these root subsystems are used to study real Del Pezzo surfaces. Since this subject is studied quite extensively it is not easy to cite all the relevant literature, but a nice overview can be found in Dolgachev [2012].

In this paper we will address the following two classifications from an algorithmic point of view:

- the classification of root subsystems of weak Del Pezzo surfaces (Du Val [1934],Dolgachev [2012]), and
- the real structures of weak Del Pezzo surfaces (Wall [1987]).

More precisely, in this paper we address to the following 2 problems.

**Problem 1.** Compute a table which classifies weak Del Pezzo surfaces up to Weyl equivalence. Provide methods which for each entry in this table enables to do explicit calculations in the Picard group of weak Del Pezzo surfaces in the corresponding equivalence class. Moreover provide algorithms to compute explicit parametrizations of weak Del Pezzo surfaces in a given equivalence class.

A solution to this problem is provided by Algorithm 11 and §6. In §6 we use methods from §5 for constructing linear series. We introduce coordinates in order to do explicit computations. Since Del Pezzo surfaces are studied so intensively it is dangerous to claim originality for every single aspect of the solution we provide. The main contribution is the algorithmic point of view and the application to weak Del Pezzo surfaces in a unified manner. Moreover we provide an implementation of the algorithms.

**Problem 2.** Compute a table which classifies conjugacy classes of real structures on weak Del Pezzo surfaces. Provide explicit coordinate representations for the action of real structures on the Picard group.

A solution to this problem is provided by Theorem 35. The method of classifying the real structures does not use the existing classification in Wall [1987] and thus provides an alternative proof. In Wall [1987] smooth Del Pezzo surfaces are considered. We consider real structures on weak Del Pezzo surfaces (which might have singularities) and do explicit computations for given singularities (see Example 36).

## 1.2 Motivation

The author needs the results of this paper for forthcoming publications concerning families of curves on surfaces (in particular families of circles).

We will recall some results which are considered well known to algebraic geometers. As a result the paper is rather lengthly; I hope this makes the material easier accessible to a larger community. For example Darboux cyclides are the projection of anticanonical models of weak degree 4 Del Pezzo surfaces. The solutions to problem 1 and problem 2 leads to classification of Darboux cyclides up to diffeomorphism. Such surfaces are of interest to applied geometers.

The algorithms for problem 1 in this paper are improvements of the algorithms in the PhD thesis: Lubbes [2011]. Since there are more than 200 cases to consider, our experience is that mistakes are easily made when computed by hand. I hope that this paper is of service to others and saves a lot of time.

## 1.3 Overview

The first part recalls some definitions concerning root systems. We also introduce non-standard definitions, namely the complement and double complement of a root system. We proof that root subsystems are equivalent if and only if their Dynkin diagram, complement and double complement are isomorphic. We introduce C1 labels, with which we will represent equivalence classes of C1 root subsystems. We present an algorithm which computes the classification of root subsystems of C1 root systems in terms of C1 labels.

In the second part we recall some theory concerning weak Del Pezzo surfaces and their Picard group. In particular we define the Del Pezzo zero-set, the Del Pezzo one-set and the Del Pezzo two-set. The first two sets are also known as the classes of respectively minus two curves and minus one curves. The classes in the Del Pezzo two-set, which cannot be written as the sum of two effective classes, are the divisor classes of one-dimensional families of conics. We say that weak Del Pezzo surfaces are Cremona equivalent if their Picard groups, with additional structure, are isomorphic. We can associate a root subsystem to a weak Del Pezzo surface; we call this a C1 object. We recall that two weak Del Pezzo surfaces are Cremona equivalent if and only if their C1 objects are isomorphic.

Next we describe an algorithm which analyses the base points of a plane linear series. We also describe an algorithm which constructs a linear series with (possibly infinitely near) base points as input. Then we show examples for constructing a parametrization of a weak Del Pezzo surface from a C1 label. In particular the C1 object of the weak Del Pezzo surface is isomorphic to the root subsystem represented by the C1 label.

Finally we present an algorithm which outputs conjugacy classes of real structures of weak Del Pezzo surfaces. The proof is independent of known classifications. As an application we consider families of conics on real weak Del Pezzo surfaces of degree four.

## Overview table of environments

| type | # | description |
|---|---|---|
| **Section** | 1 | Introduction |
| **Subsection** | 1.1 | Problems |
| **Subsection** | 1.2 | Motivation |
| **Subsection** | 1.3 | Overview |
| **Subsection** | 1.4 | Notation |
| **Section** | 2 | Root subsystems |
| **Subsection** | 2.1 | Definitions |
| Definition | 1 | root (sub)system and the Weyl group |
| Definition | 2 | category of root systems and root subsystems |
| Definition | 3 | attributes of roots |
| Definition | 4 | Cartan matrix and Dynkin diagram |
| **Subsection** | 2.2 | C1 labels |
| Definition | 5 | C1 root system and C1 category |
| Definition | 6 | C1 kit |
| Proposition | 7 | properties of C1 kit |
| Definition | 8 | C1 label |
| **Subsection** | 2.3 | Complement |
| Definition | 9 | complement and double complement of root subsystems |
| Proposition | 10 | properties of complement and double complement of root subsystems |
| **Section** | 3 | Algorithm: geometric bases of C1 root subsystems |
| Algorithm | 11 | geometric bases of C1 root subsystems |
| Proposition | 12 | geometric bases of C1 root subsystems |
| **Section** | 4 | Weak Del Pezzo surfaces |
| **Subsection** | 4.1 | Definitions |
| Definition | 13 | weak Del Pezzo surface |
| Definition | 14 | degree of weak Del Pezzo surface |
| Definition | 15 | standard Del Pezzo and geometrically ruled basis |
| Proposition | 16 | properties of standard Del Pezzo basis |
| **Subsection** | 4.2 | Del Pezzo sets |
| Definition | 17 | indecomposable class |
| Definition | 18 | Del Pezzo zero-set |
| Definition | 19 | Del Pezzo one-set |
| Definition | 20 | Del Pezzo two-set |
| Proposition | 21 | properties of Del Pezzo sets and intersection product |
| **Subsection** | 4.3 | Cremona equivalence |
| Definition | 22 | enhanced Picard group |
| Definition | 23 | Cremona equivalence of weak Del Pezzo surfaces |
| Definition | 24 | C1 object of weak Del Pezzo surface |

| | | |
|---|---|---|
| Proposition | 25 | properties of Cremona equivalence of weak Del Pezzo surfaces |
| **Section** | 5 | Algorithm: analyzing and constructing linear series |
| Example | 26 | base point analysis of linear series |
| Example | 27 | linear series with prescribed base points |
| Algorithm | 28 | base point analysis of linear series in the projective plane |
| Algorithm | 29 | create linear series in the projective plane with given base points |
| **Section** | 6 | Examples of constructing weak Del Pezzo surfaces |
| Example | 30 | degree four Del Pezzo with two ordinary double points |
| Example | 31 | degree one Del Pezzo with four cusps |
| **Section** | 7 | Algorithm: real structures of weak Del Pezzo surfaces |
| Definition | 32 | real structure for weak Del Pezzo surfaces |
| Algorithm | 33 | real structures of weak Del Pezzo surfaces |
| Example | 34 | real structures of weak Del Pezzo surfaces |
| Theorem | 35 | real structures of weak Del Pezzo surfaces |
| Example | 36 | real structures of weak Del Pezzo surface of degree four |
| **Section** | 8 | Acknowledgements |
| **Appendix** | A | Table of geometric C1 labels |
| Table | 37 | geometric bases of C1 root subsystems up to Weyl equivalence |
| **Appendix** | B | Real structures of weak Del Pezzo surfaces |
| Table | 38 | real structures of weak Del Pezzo surfaces |

## 1.4 Notation

Let $V$ be an Euclidean space with bilinear form $(\cdot,\cdot)$. We also assume a total order $>$ on $V$. Let $\check{v} = \frac{2v}{(v,v)}$ for all $v$ in $V - \{0\}$. The reflection along a hyperplane which is orthogonal to $u \in V - \{0\}$ is defined by $\tau_u : V \to V, \quad v \mapsto v - (\check{u}, v)u$.

Later in §5 we will denote an algebraically closed number field by $\mathbf{F}$. First occuring in §4, the map associated to a divisor class $D$ is denoted by $\varphi_D$.

# 2 Root subsystems

We recall some theory concerning root subsystems, which can be found in for example Bourbaki [2002] and de Graaf [2000].

## 2.1 Definitions

**Definition 1. (root (sub)system and the Weyl group)** Let $R \subset V - \{0\}$ be a finite subset. We call $R$ a *root system* in $V$ if and only if $V = \langle R \rangle$, $\tau_r(s) \in R$, $(\check{r}, s) \in \mathbf{Z}$, and $\mathbf{R}\langle r \rangle \cap R = \{\pm r\}$, for all $r, s \in R$. We call $S \subset R$ a *root subsystem* of root system $R$ if and only if $S$ is a root system in the Euclidean vector subspace $\mathbf{R}\langle S \rangle \subset V$. The *Weyl group* of a root system $R$ is defined as $W(R) = \langle\ \tau_r \ \mid\ r \in R\ \rangle$ where multiplication of reflections is defined as composition.

**Definition 2. (category of root systems and root subsystems)** The *category of root systems* $\mathcal{R}$ is defined as follows:

- The objects of $\mathcal{R}$ are root systems.
- The morphisms of $\mathcal{R}$ are $(R \subset V) \xrightarrow{f} (R' \subset V')$ such that $V \xrightarrow{f} V'$ is an Euclidean isometry and $f(R) = R'$.

Let $\mathcal{R}$ be the category of root systems. Let $\mathcal{R}'$ be the skeleton category of $\mathcal{R}$ (each object in $\mathcal{R}'$ uniquely represents an isomorphism class of an object in $\mathcal{R}$). The *category of root subsystems* $\mathcal{S}$ is defined as follows:

- The objects of $\mathcal{S}$ are pairs $(S, R)$ where $R \in \mathrm{Obj}\mathcal{R}'$ and $S \subset R$ is a root subsystem.
- The morphisms of $\mathcal{S}$ are $(S, R) \xrightarrow{f} (S', R)$ such that $R \xrightarrow{f} R$ is an element of the Weyl group acting on $R$ and $f(S) = S'$.

**Definition 3. (attributes of roots)** Let $R$ be a root system in the Euclidean vector space $V$ with total order $>$. We call $r \in R$ *positive* if and only if $r > 0$. We call $r \in R$ *simple* if and only if $r$ is positive and there don't exists positive roots $s, t$ such that $r = s + t$. We call $r \in R$ *maximal* if and only if $r > s$ or $r = s$ for all $s \in R$.

**Definition 4. (Cartan matrix and Dynkin diagram)** Let $R$ be a root system in the Euclidean vector space $V$ with total order $>$. Let $B = (b_i)_{i \in I}$ be the indexed set of simple roots. The *Cartan matrix* of $B$ is defined as $M = (m_{ij})_{i,j \in I}$ where $m_{ij} = (\check{b_i}, b_j)$ for all $i, j \in I$. The *Dynkin diagram* $D(R)$ of $R$ is defined as a graph where

- the vertices are labeled by the simple roots $b_i$,
- $b_i$ and $b_j$ are connected by $m_{ij}m_{ji}$ lines and
- an arrow pointing from the larger to the smaller vector if $m_{ij}m_{ji} > 1$.

for all $i, j \in I$.

We recall that the set of simple roots of a root system form a *basis* for a root system.

## 2.2 C1 labels

**Definition 5. (C1 root system and C1 category)** We call a root system $R$ a *C1 root system* if and only if $D(R)$ is of either one of the following types $A_1$, $A_1+A_2$, $A_4$, $D_5$, $E_6$, $E_7$ or $E_8$. Let $\mathcal{S}$ be the category of root subsystems. The *C1 category* $\mathcal{Z}$ is defined as the largest subcategory of $\mathcal{S}$ such that $R$ is a C1 root system for all $(S,R) \in \mathrm{Obj}\mathcal{Z}$. The name C1 has some 'historical' reasons with C standing for Cartan. However, I have to admit that at this point the naming is bit arbitrary, and now it is too late to change.

**Definition 6. (C1 kit)** A *C1 kit* is defined as

$$(M, (\cdot,\cdot), <, R)$$

where

- $M = \mathbf{Z}\langle H, Q_1, \ldots, Q_r\rangle$ is a $\mathbf{Z}$-module with basis and an element of $M$ is often given by its coefficient vector with respect to this basis,
- $r \in [2, \ldots, 8]$,
- $(\cdot,\cdot)$ is the bilinear form on $M \underset{\mathbf{Z}}{\otimes} \mathbf{R}$ with signature $(+,-,\ldots,-)$,
- $m > m'$ if and only if $(m_0, -m_1, \ldots, -m_r) >_{lex} (m'_0, -m'_1, \ldots, -m'_r)$ for all $m, m' \in M$, and
- $R = \{\ m \in M \ \mid\ (m,k) = 0 \text{ and } (m,m) = -2\ \}$ for $k = (-3, 1, \ldots, 1)$.

**Proposition 7. (properties of C1 kit)**

Let $(M, (\cdot,\cdot), <, R)$ be a C1 kit. Let $V = \{\ v \in M \underset{\mathbf{Z}}{\otimes} \mathbf{R} \ \mid\ (v,k) = 0\ \}$ where $k = (-3, 1, \ldots, 1)$.

We have that $R$ is a C1 root system in the vector space $V$ and

| rank of $M$ | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|---|---|---|---|---|---|---|---|
| $D(R)$ | $A_1$ | $A_1 + A_2$ | $A_4$ | $D_5$ | $E_6$ | $E_7$ | $E_8$ |

where $D(R)$ is the Dynkin diagram of $R$.

**Proof:** This is well known: Manin [1986]. ☕

**Definition 8. (C1 label)** The set of *C1 label elements* is defined as $G = \{\ \pm ab, \pm 1abc, \pm 2ab, \pm 30a \ \mid\ a,b,c \in [1,\ldots,8]\ \}$, A *C1 label* is defined as $(L, r)$ where $L \subset G$ and $r \in [1, \ldots, 8]$. Let $(M, (\cdot,\cdot), <, R)$ be a C1 kit. The *C1 label elements to root function* $G \overset{\varphi}{\to} R$ is defined as follows:

- $\varphi(ab) = Q_a - Q_b$,
- $\varphi(1abc) = H - Q_a - Q_b - Q_c$,
- $\varphi(2ab) = 2H - \sum\limits_{i \in [1,\ldots,8], i \notin \{a,b\}} Q_i$, and

- $\varphi(30a) = 3H - \sum_{i \in [1,\ldots,8], i \notin \{a\}} Q_i - 2Q_a$,

for all $a, b, c \in [1, \ldots, 8]$ and $\varphi(-l) = -\varphi(l)$ for all $l \in G$. We call a C1 label *geometric* if and only if the label does not contain minus symbols.

We will use C1 labels $(L, r)$ for representing a basis $(\ \varphi(l)\ )_{l \in L}$ of a root subsystem $S$ of a C1 root system $R$. In other words we can represent isomorphism classes of root subsystems by C1 labels.

## 2.3 Complement

The following definition and proposition is non-standard and used for Algorithm 11.

**Definition 9. (complement and double complement of root subsystems)** Let $R$ be a root system in $V$. Let $S \subset R$ be a root subsystem. The *complement root subsystem* is defined as $C_R(S) = \{\ r \in R \ \mid\ (r, s) = 0 \text{ for all } s \in S\ \}$. The *double complement root subsystem* is defined as $K_R(S) = \{\ r \in R \ \mid\ r \in \mathbf{R}\langle S \rangle\ \}$.

**Proposition 10. (properties of complement and double complement of root subsystems)**

Let $\mathcal{Z}$ be the category of C1 root subsystems. Let $(S, R)$ and $(S', R)$ in Obj$\mathcal{Z}$ such that $R$ is irreducible. Let $D(S)$ denote the Dynkin diagram of $S$.

We have that $(S, R) \cong (S', R)$ if and only if $D(S) \cong D(S')$, $|C_R(S)| = |C_R(S')|$ and $|K_R(S)| = |K_R(S')|$.

Here $\cong$ are isomorphisms in the category of respectively root subsystems and graphs.

**Proof:**

*Claim:* We have that $\Longrightarrow$.
We have that $C_R(S) = \mathbf{R}\langle S \rangle^{\perp} \cap R$ where $\mathbf{R}\langle S \rangle \subset V$ is an Euclidean vector subspace. Similarly we have that $K_R(S) = \mathbf{R}\langle S \rangle^{\perp\perp} \cap R$. We have an action $R \xrightarrow{f} R$ of an element the Weyl group such that $f(S) = S'$. This extends linearly to an Euclidean isometry $V \xrightarrow{f'} V$ such that $f'(R) = R$. From $f'$ preserving orthogonality it follows that this claim holds.

*Claim:* We have that $\Longleftarrow$.
This claim follows from tables 10.1 and 10.2 in Oshima [2006]. These tables contain $D(C_R(S))$ for each root subsystem $S$ of C1 root systems. We find that if $D(S) \cong D(S')$, $C_R(S) \cong C_R(S')$ then $(S, R) \cong (S', R)$, for almost all $S, S'$. Except for the subsystems $4A_1$, $A_5 + A_1$ and $A_3 + 2A_1$ in $E_7$ we see that there are two Weyl equivalence classes which cannot be distinguished this way. For example there exists root subsystems $S$ and $S'$ of root system $R$ such that $D(R) = E_7$, $D(S) = D(S') = A_5 + A_1$ and $D(C_R(S)) = D(C_R(S')) = \emptyset$. However for these cases we find that $K_R(S) \ncong_b K_R(S')$ and thus this claim holds. ☕

# 3 Algorithm: geometric bases of C1 root subsystems

The following algorithm computes geometric C1 labels for C1 root subsystems. Later we will use the output to construct explicit examples, and to do computations in the Picard group of weak Del Pezzo surfaces. The classification of C1 root subsystems is well known (see section 8.2.3 in Dolgachev [2012]).

**Algorithm 11. (geometric bases of C1 root subsystems)**

**Output:** A list of geometric C1 labels (thus without minus symbols). Each geometric C1 label represents a basis for a unique representative of each isomorphism class in the category of C1 root subsystems. Recall that isomorphisms of root subsystems are defined by actions of the Weyl group.

- Let $G \overset{\varphi}{\to} R$ be the C1 label elements to root function. By abuse of notation we will not distinguish between a C1 label $(L, r)$, and the corresponding root set $(\ \varphi(l)\ )_{l\in L}$. For example the bilinear form of C1 labels $L_i$ is defined by the bilinear form of $(\ \varphi(l)\ )_{l\in L_i}$. Similar for linear independence. For representing roots internally we use the C1 kit.
- Let
  $Z_0 := \{\ 12,\ 23,\ 34,\ 45,\ 56,\ 67,\ 78,\ 1123,\ -1145,\ -1345,\ -1167,\ -1178,\ -278,\ -218,\ -308,\ 218\ \}$,
  $Z_1 := \{\ 1123,\ 12,\ 23,\ 34,\ 45,\ 56,\ 67,\ 78\ \}$,
  $Z_2 := \{\ 1123,\ 12,\ 23,\ 34,\ 45,\ 56,\ 67,\ 78,\ 1145,\ 1347,\ 1678,\ 1127,\ 1456,\ 1567,\ 234,\ 278,\ 308\ \}$,
  $Z_3 := \{\ 1123,\ 1345,\ 1165,\ 1285,\ 1673,\ 1274,\ 1684,\ 1178\ \}$.
- We apply the following two steps by looping over $r \in [2, \ldots, 8]$ and $i \in [0, \ldots, 3]$.
- Let $Y_i$ be the set of C1 labels $(L, r)$ where $L$ is a subset of $Z_i \cap \{\ \pm ab, \pm 1abc, \pm 2ab, \pm 30a \mid a, b, c \in [r]\ \}$ such that $L$ is linear independent and $(l, m) \in \{0, 1\}$ for all $l, m \in L$. So $L$ defines a valid basis for a root subsystem of rank $r$.
- Using Proposition 10 we reduce the set $Y_i$ such that it only contains non-isomorphic root subsystems represented by C1 labels $(L, r)$. Sage contains a function which checks isomorphisms of graphs. We check for the following special case: the C1 labels $(\{1123\}, 3)$ and $(\{23\}, 3)$ are non-isomorphic root systems of Dynkin type $A_1$ in the reducible (!) root system of type $A_1 + A_2$ (although their complement and double complement root subsystems are isomorphic).
- We have that $\cup_i Y_i$ contains isomorphic root subsystems. For a given equivalence class of root subsytems in $F_0$ the preferred representative is in $F_1$. If it does not exists in $F_1$ then $F_2$. If it does not exists in $F_2$ then $F_3$. If it does not exists in $F_3$ then $F_0$.
- We return the obtained set of C1 labels.

**Proposition 12. (geometric bases of C1 root subsystems)**

Algorithm 11 is correct.

**Proof:** The set $Z_0$ is constructed using Dynkin's Algorithm, which recursively adds the negative of maximal roots of sub diagrams. This ensures independently of existing classifications that all isomorphism classes of root subsystems is reached in $Z_0$. The set $Z_i$ for $i > 0$ are contructed such that each equivalence class of root subsystems has a geometric C1 label. The set $Z_1$ forms a basis for the root system of Dynkin type $E_8$. If we apply the algorithm using only $Z_0$ and $Z_1$ then the output contains C1 labels which are not geometric. We construct sets $Z_2$ and $Z_3$ such that each of these non-geometric C1 labels is represented by a geometric C1 label. The root subsystem $4A_2$ in $E_8$ requires extra attention (see Example 31) and leads to the labels of $Z_3$. From Proposition 10 it follows that no two C1 labels in in $T$ represent isomorphic root subsystems of an irreducible root system. When the rank is 3 then we consider root subsystems of a reducible root system $A_1 + A_2$. We have to make an explicit check for two non-isomorphic root subsystems of Dynkin type $A_1$. It can be verified by known classifications (see for example Dolgachev [2012]) that each isomorphism class of root subsystems is represented by an element of $T$. The author would like to thank Ulrich Derenthal for pointing out the missing $A_1$ case when the rank equals 3. ☕

The output of Algorithm 11 is in Table 37.

In order to motivate the following sections in relation with the previously described algorithm we give a flash forward before recalling some definitions of weak Del Pezzo surfaces.

In §5 we will recall how to create a linear series in the projective plane with a prescribed set of base points. A linear series of degree 3 in the projective plane, with for example 5 generic base points defines a parametrization of the anticanonical model of a weak Del Pezzo surface X of degree 4. Now if three base points lie on a line, and the remaining base points are generic then this corresponds to an effective minus two class. The Picard group of X together with Betti numbers is determined by the set of effective minus two classes. If we assign indices to the 5 base points $p_1, \ldots, p_5$, then the C1 label $(\{1123\}, 5)$ denotes that the points $p_1$, $p_2$ and $p_3$ lie on a line and that $p_4$ and $p_5$ are in generic position. In Example 30 we will the result for the C1 label $(\{1123, 45\}, 5)$.

We will define the enhanced Picard group of X to be the Picard group together with intersection product, Betti numbers and the anticanonical divisor. We recall that enhanced Picard groups of weak Del Pezzo surfaces are isomorphic if and only if the root subsystem associated to the effective minus two classes are isomorphic.

Finally we consider an additional involution on the Picard group, which corresponds to a real structure. We will propose an algorithm to find up to ismorphism of real enhanced Picard groups an explicit description of the real structures. As an application we classify in Example 36 the real families of conics of anticanonical models of real weak Del Pezzo surface of degree 4.

# 4 Weak Del Pezzo surfaces

In this section we will recall some theory of Del Pezzo surfaces and its Picard group. References for this section are Manin [1986] and Dolgachev [2012]. Details can be found in appendix E of Lubbes [2011].

## 4.1 Definitions

**Definition 13. (weak Del Pezzo surface)** Let X be a nonsingular complex projective surface. Let $K$ be the canonical class of X. We call X a *weak Del Pezzo surface* if and only if $-K$ is nef and big.

**Definition 14. (degree of weak Del Pezzo surface)** Let X be a weak Del Pezzo surface. Let $K$ be the canonical class of X. The *degree* of X is defined as $K^2$.

**Definition 15. (standard Del Pezzo and geometrically ruled basis)** A *standard Del Pezzo basis* $B(r)$ is defined as a unimodular lattice

$$B(r) = \mathbf{Z}\langle\ H,\ Q_1,\ \ldots,\ Q_r\ \rangle,$$

with bilinear form $H^2 = 1$, $Q_iQ_j = -\delta_{ij}$ and $HQ_i = 0$. A *standard geometrically ruled basis* $P(r)$ is defined as a unimodular lattice

$$P(r) = \mathbf{Z}\langle H, F\rangle$$

with bilinear form $F^2 = 0$, $FH = 1$ and $H^2 = r$.

**Proposition 16. (properties of standard Del Pezzo basis)**

Let X be a weak Del Pezzo surface. Let $K$ be the canonical class of X. Let $B(r) = \mathbf{Z}\langle\ H,\ Q_1,\ \ldots,\ Q_r\ \rangle$ be the standard Del Pezzo basis. Let $A(\mathrm{X})$ be the Picard group of X together with intersection product and the element $K$.

**a)** We have that $1 \leq K^2 \leq 9$.

**b)** If $K^2 \neq 8$ then $A(\mathrm{X}) \cong B(9 - K^2)$ with $-K = 3H - Q_1 - \ldots - Q_r$.

Let $P(r) = \mathbf{Z}\langle\ H,\ F\ \rangle$ be the standard geometrically ruled basis.

**c)** If $K^2 = 8$ then $A(\mathrm{X}) = P(r)$ with $-K = 2H - (r-2)F$ for $r = 0, 1$ or $2$.

**Proof:** See appendix E, section 2 of Lubbes [2011] or Dolgachev [2012]. ☕

## 4.2 Del Pezzo sets

We state properties of divisor classes of weak Del Pezzo surfaces. We distinguish between three different sets of divisor classes. The *Del Pezzo zero-set* are the minus two classes (which have intersection zero with the anticanonical class). The *Del Pezzo one-set* are the minus one classes (which have intersection one with the anticanonical class). And the *Del Pezzo two-set* are the divisor classes with self-intersection zero and intersection two with the anticanonical class.

We recall that the indecomposable fixed classes of weak Del Pezzo surfaces are either minus one or minus two classes. We state some properties concerning the decomposition of divisor classes in Del Pezzo sets.

We recall that classes in the Del Pezzo two-set are one dimensional rational linear series. If this linear series has fixed components then these components must be effective minus two classes. The linear series defined by the Del Pezzo two-set define 1 dimensional family of conics.

The classes in the indecomposable Del Pezzo zero-set determine the singularities of the anticanonical model of a weak Del Pezzo surface. The indecomposable Del Pezzo one-set defines the lines of the anticanonical model.

Later, when we will introduce real structures on the Picard group, we will see that the tuple defined by the number of classes in the Del Pezzo zero-, one- and two-set, which are fixed under the real structure, are strong invariants for the real structure.

**Definition 17. (indecomposable class)** Let $X$ be a nonsingular complex projective variety. We call $D \in \text{PicX}$ an *indecomposable class* if and only if $D$ is effective and can not be written as a positive sum of two effective classes.

**Definition 18. (Del Pezzo zero-set)** Let X be a weak Del Pezzo surface. Let $K$ be the canonical divisor class of $X$. Let $B(r) = \mathbf{Z}\langle\ H,\ Q_1,\ \ldots,\ Q_r\ \rangle$ be the standard Del Pezzo basis. Let $P(r) = \mathbf{Z}\langle\ H,\ F\ \rangle$ be the standard geometrically ruled basis. The *Del Pezzo zero-set* $F(\text{X})$ is defined as the set of all classes in Pic(X) which, up to permutation of the $Q_i$, have coefficients as in the following table:

| $K^2$ | $H$ | $Q_1$ | $Q_2$ | $Q_3$ | $Q_4$ | $Q_5$ | $Q_6$ | $Q_7$ | $Q_8$ |
|---|---|---|---|---|---|---|---|---|---|
| $\leq 7$ |  | 1 | $-1$ |  |  |  |  |  |  |
| $\leq 6$ | 1 | $-1$ | $-1$ | $-1$ |  |  |  |  |  |
| $\leq 3$ | 2 | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ |  |  |
| 1 | 3 | $-2$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ |

If $K^2 = 8$ and $\text{Pic(X)} = P(0)$ then $F(\text{X}) = \{\pm H - F\}$. If $K^2 = 8$ and $\text{Pic(X)} = P(1)$ then $F(\text{X}) = \emptyset$. If $K^2 = 8$ and $\text{Pic(X)} = P(2)$ then $F(\text{X}) = \{\pm H - 2F\}$. The *indecomposable Del Pezzo zero-set* is defined as

$$F_{\geq 0}(\text{X}) = \{\ C \in F(\text{X})\ \ |\ \ C \text{ is an indecomposable class }\}.$$

**Definition 19. (Del Pezzo one-set)** Let X be a weak Del Pezzo surface. Let $K$ be the canonical divisor class of $X$. Let $B(r) = \mathbf{Z}\langle\ H,\ Q_1,\ \ldots,\ Q_r\ \rangle$ be the standard Del Pezzo basis. Let $P(r) = \mathbf{Z}\langle\ H,\ F\ \rangle$ be the standard geometrically ruled basis. The *Del Pezzo one-set* $E(\text{X})$ is defined as the set of all classes in Pic(X) which, up to permutation of the $Q_i$, have coefficients as in the following table:

| $K^2$ | $H$ | $Q_1$ | $Q_2$ | $Q_3$ | $Q_4$ | $Q_5$ | $Q_6$ | $Q_7$ | $Q_8$ |
|---|---|---|---|---|---|---|---|---|---|
| $\leq 7$ |  | 1 |  |  |  |  |  |  |  |
| $\leq 7$ | 1 | $-1$ | $-1$ |  |  |  |  |  |  |
| $\leq 4$ | 2 | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ |  |  |  |
| $\leq 2$ | 3 | $-2$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ |  |
| 1 | 4 | $-2$ | $-2$ | $-2$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ |
| 1 | 5 | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-1$ | $-1$ |
| 1 | 6 | $-3$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ |

If $K^2 = 8$ and $\mathrm{Pic}(\mathrm{X}) = P(0)$ then $F(\mathrm{X}) = \emptyset$. If $K^2 = 8$ and $\mathrm{Pic}(\mathrm{X}) = P(1)$ then $F(\mathrm{X}) = \{H - F\}$. If $K^2 = 8$ and $\mathrm{Pic}(\mathrm{X}) = P(2)$ then $F(\mathrm{X}) = \emptyset$. The *indecomposable Del Pezzo one-set* is defined as

$$E_{\geq 0}(\mathrm{X}) = \{ \ C \in E(\mathrm{X}) \ \mid \ C \text{ is an indecomposable class } \}.$$

**Definition 20. (Del Pezzo two-set)** Let X be a weak Del Pezzo surface. Let $K$ be the canonical divisor class of $X$. Let $B(r) = \mathbf{Z}\langle \ H, \ Q_1, \ \ldots, \ Q_r \ \rangle$ be the standard Del Pezzo basis. Let $P(r) = \mathbf{Z}\langle \ H, \ F \ \rangle$ be the standard geometrically ruled basis. The *Del Pezzo two-set* $G(\mathrm{X})$ is defined as the set of classes in $\mathrm{Pic}(\mathrm{X})$ which, up to permutation of the $Q_i$, have coefficients as in the following table:

| $K^2$ | $H$ | $Q_1$ | $Q_2$ | $Q_3$ | $Q_4$ | $Q_5$ | $Q_6$ | $Q_7$ | $Q_8$ |
|---|---|---|---|---|---|---|---|---|---|
| $\leq 7$ | 1 | $-1$ | | | | | | | |
| $\leq 5$ | 2 | $-1$ | $-1$ | $-1$ | $-1$ | | | | |
| $\leq 3$ | 3 | $-2$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | | |
| $\leq 2$ | 4 | $-2$ | $-2$ | $-2$ | $-1$ | $-1$ | $-1$ | $-1$ | |
| $\leq 2$ | 5 | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-1$ | |
| 1 | 4 | $-3$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ | $-1$ |
| 1 | 5 | $-3$ | $-2$ | $-2$ | $-2$ | $-1$ | $-1$ | $-1$ | $-1$ |
| 1 | 6 | $-3$ | $-3$ | $-2$ | $-2$ | $-2$ | $-2$ | $-1$ | $-1$ |
| 1 | 7 | $-3$ | $-3$ | $-3$ | $-3$ | $-2$ | $-2$ | $-2$ | $-1$ |
| 1 | 7 | $-4$ | $-3$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ | $-2$ |
| 1 | 8 | $-3$ | $-3$ | $-3$ | $-3$ | $-3$ | $-3$ | $-3$ | $-1$ |
| 1 | 8 | $-4$ | $-3$ | $-3$ | $-3$ | $-3$ | $-2$ | $-2$ | $-2$ |
| 1 | 9 | $-4$ | $-4$ | $-3$ | $-3$ | $-3$ | $-3$ | $-3$ | $-2$ |
| 1 | 10 | $-4$ | $-4$ | $-4$ | $-4$ | $-3$ | $-3$ | $-3$ | $-3$ |
| 1 | 11 | $-4$ | $-4$ | $-4$ | $-4$ | $-4$ | $-4$ | $-4$ | $-3$ |

If $K^2 = 8$ and $\mathrm{Pic}(\mathrm{X}) = P(0)$ then $F(\mathrm{X}) = \{H\}$. If $K^2 = 8$ and $\mathrm{Pic}(\mathrm{X}) = P(1)$ then $F(\mathrm{X}) = \{F\}$. If $K^2 = 8$ and $\mathrm{Pic}(\mathrm{X}) = P(2)$ then $F(\mathrm{X}) = \{H - F\}$. The *irreducible Del Pezzo two-set* is defined as

$$G_{irr}(\mathrm{X}) = \{ \ C \in G(\mathrm{X}) \ \mid \ \text{the linear series } |C| \text{ has no fixed components } \}.$$

**Proposition 21. (properties of Del Pezzo sets and intersection product)** Let X be a weak Del Pezzo surface. Let $K$ be the canonical divisor class of X. Let $F(\mathrm{X})$ be the Del Pezzo zero-set. Let $E(\mathrm{X})$ be the Del Pezzo one-set. Let $G(\mathrm{X})$ be the Del Pezzo two-set.

**a)** We have that

- $F(\mathrm{X}) = \{ \ C \in \mathrm{Pic}(\mathrm{X}) \ \mid \ -KC = 0 \text{ and } C^2 = -2 \ \}$,
- $E(\mathrm{X}) = \{ \ C \in \mathrm{Pic}(\mathrm{X}) \ \mid \ -KC = 1 \text{ and } C^2 = -1 \ \}$, and
- $G(\mathrm{X}) = \{ \ C \in \mathrm{Pic}(\mathrm{X}) \ \mid \ -KC = 2 \text{ and } C^2 = 0 \ \}$.

**b)** We have that $E_{\geq 0} = \{ \ C \in \mathrm{Pic}(\mathrm{X}) \ \mid \ C \in E(\mathrm{X}) \text{ such that } CF \geq 0 \text{ for all } F \in F_{\geq 0}(\mathrm{X}) \ \}$.

**c)** We have that $G_{irr} = \{ \ C \in \mathrm{Pic}(\mathrm{X}) \ \mid \ C \in G(\mathrm{X}) \text{ such that } CF \geq 0 \text{ for all } F \in F_{\geq 0}(\mathrm{X}) \ \}$.

**Proof:** For **a)** see appendix E, section 3, proposition 223 of Lubbes [2011]. For **b)** see section 8.2.6, Lemma 8.2.18 in Dolgachev [2012]. For **c)** see appendix E, section 3, proposition 226 of Lubbes [2011]. ☕

## 4.3 Cremona equivalence

The anticanonical model Y of a weak Del Pezzo surface X is defined as $\mathrm{X} \stackrel{\varphi_D}{\to} \mathrm{Y} \subset \mathbf{P}^n$ where $D = -mK$ for some positive integer $m$ such that $\varphi_D$ is birational.

The indecomposable Del Pezzo zero-set $F_{\geq 0}(\mathrm{X})$ determines the singularities of Y. Moreover, we recall that the effective minus two classes in $F_{\geq 0}(\mathrm{X})$ determine the Picard group together with intersection product, distinguished anticanonical class and Betti numbers.

The Del Pezzo zero-set $F(\mathrm{X})$ defines a C1 root system $R$ in the vector space $\mathrm{Pic}\mathrm{X} \otimes \mathbf{R}$. We have that $F_{\geq 0}(\mathrm{X})$ defines a root subsystem of $R$. So we can associate a root subsystem to a weak Del Pezzo surface.

We consider two weak Del Pezzo surfaces to be *Cremona equivalent* if and only if their associated root subsystems are isomorphic.

See appendix E, section 4 in Lubbes [2011] or 8.2.8, proposition 24 in Dolgachev [2012].

**Definition 22. (enhanced Picard group)** Let X a weak Del Pezzo surface. Let $K$ be the canonical divisor class on X. The *enhanced Picard group* $A(\mathrm{X})$ is defined as

$$(\ \mathrm{Pic}(\mathrm{X}),\ K,\ \cdot,\ h\ )$$

where $\mathrm{Pic}(\mathrm{X})$ is the Picard group, $\mathrm{Pic}(\mathrm{X}) \times \mathrm{Pic}(\mathrm{X}) \stackrel{\cdot}{\to} \mathbf{Z}$ is the intersection product on divisor classes, and $\mathbf{Z} \times \mathrm{Pic}(\mathrm{X}) \stackrel{h}{\to} \mathbf{Z}_{\geq 0}$ assigns the $i$-th Betti number to a divisor class for $i \in \mathbf{Z}$. For $h(i, D)$ we use the notation $h^i(D)$.

**Definition 23. (Cremona equivalence of weak Del Pezzo surfaces)** Let X and X′ be weak Del Pezzo surfaces. Let $A(\mathrm{X})$ be the enhanced Picard group. The *Cremona equivalence relation of weak Del Pezzo surfaces* is defined as follows:

$$\mathrm{X} \sim_w \mathrm{X}' \quad :\Leftrightarrow \quad A(\mathrm{X}) \cong A(\mathrm{X}').$$

**Definition 24. (C1 object of weak Del Pezzo surface)** Let $F(\mathrm{X})$ be the Del Pezzo zero-set. Let $\mathcal{Z}$ be the C1 category. The *C1 object* of X is defined as

$$W(\mathrm{X}) := (S, R) \in \mathrm{Obj}\mathcal{Z}$$

where $R$ is the root system defined by $F(\mathrm{X})$, and $S$ is the root subsystem of $R$ defined by $F_{\geq 0}(\mathrm{X})$. See Proposition 7.

**Proposition 25. (properties of Cremona equivalence of weak Del Pezzo surfaces)** Let X and X′ be weak Del Pezzo surfaces. Let $A(\mathrm{X})$ be the enhanced Picard group. Let $W(\mathrm{X})$ be the C1 object.

**a)** We have that $A(\mathrm{X}) \cong A(\mathrm{X}')$ if and only if $W(\mathrm{X}) \cong W(\mathrm{X}')$.

**Proof:** See appendix E, section 4 in Lubbes [2011], or 8.2.8, proposition 24 in Dolgachev [2012]. ☕

# 5 Algorithm: analyzing and constructing linear series

We recall the analysis of base points of linear series.

Let $\langle f_1, \ldots, f_n \rangle$ be a linear series such that $f_i \in \mathbf{F}[x, y, z]$ are homogeneous. Let $\mathrm{F} : F(x, y, z) = \sum\limits_{i\in[n]} c_i f_i(x, y, z)$ for $c_i \in \mathbf{F}$ for all $i \in [n]$. Let $\tilde{\mathrm{F}} \xrightarrow{\pi} \mathrm{F}$ be the blowup at the origin $p = (0 : 0 : 1)$

We consider a diagram of charts of the blowup $\pi$ in $\mathbf{P}^2 \times \mathbf{P}^1 \to \mathbf{P}^2$ with coordinates $(x : y : z) \times (s : t)$:

$$\begin{tikzcd} \tilde{U}'_{zt} \arrow[r, "\varphi"] & \tilde{U}_{zt} \arrow[r, "\pi|\tilde{U}_{zt}"] & U_z \end{tikzcd}$$

where

- $U_z = \{\ (\frac{x}{z}, \frac{y}{z})\ \mid\ F(\frac{x}{z}, \frac{y}{z}, 1) = 0 \text{ and } z \neq 0\ \}$,
- $\tilde{U}_{zt} = \{\ (\frac{x}{z}, \frac{y}{z}) \times (\frac{s}{t})\ \mid\ \frac{x}{z} = \frac{y}{z}\frac{s}{t} \text{ and } F(\frac{x}{z}, \frac{y}{z}, 1) = 0\ \} \subset \mathbf{A}^2 \times \mathbf{A}^1$,
- $\tilde{U}'_{zt} = \{\ (\frac{s}{t}, \frac{y}{z})\ \mid\ F(\frac{y}{z}\frac{s}{t}, \frac{y}{z}, 1) = 0\ \} \subset \mathbf{A}^2$, and
- $\varphi : \tilde{U}'_{zt} \to \tilde{U}_{zt}, \quad (\frac{s}{t}, \frac{y}{z}) \mapsto (\frac{s}{t}\frac{y}{z}, \frac{y}{z}) \times (\frac{s}{t})$.

We have that $\varphi$ is an isomorphism and

$$\varphi \circ \pi|\tilde{U}_{zt} : \tilde{U}'_{zt} \to U_z, \quad (u, v) \mapsto (uv, v).$$

We have that $\tilde{U}_{zt}$ is a chart of the pullback $\pi^*\mathrm{F} = \tilde{\mathrm{F}} + mE$ where in this chart we have that $E : v = 0$. From $\varphi$ being an isomorphism it follows that $\tilde{U}'_{zt} : F(uv, v, 1) = 0 \subset \mathbf{A}^2$ has a factor $v^m$ such that $m = \mathrm{mult}_p\mathrm{F}$ is the multiplicity of F at $p = (0 : 0 : 1)$.

We consider $v^{-m}F(uv, v, 1) = 0$ as our new chart of linear series. We translate the base point we want to analyze to the origin. This will be our new $U_z$, and we can repeat the procedure.

This leads to an algorithm which analyses simple and infinitely near base points. We have to keep in mind that charts may overlap. For example we first analyze the chart $U_z \subset \mathrm{F}$. Then if a base point $q$ in the chart $U_y$ has non-zero coordinate $\frac{z}{y}$ then $q \in U_z \cap U_y$. and thus we already analyzed the base point in $U_z$. Similarly for the charts $\tilde{U}_{zt}$ and $\tilde{U}_{zs}$. See chapter 8 in Lubbes [2011] for the algorithms in pseudo code.

**Example 26. (base point analysis of linear series)** We use the same notation as above. Let $\mathrm{F} : F(x, y, z) = c_0(y^2 - 2yz + z^2) + c_1xz + c_2xy + c_3x^2 = 0 \subset \mathbf{P}^2$ be a linear series with $(c_0 : \ldots : c_3) \in \mathbf{P}^3$.

We find that $U_z : c_0(v^2 - 2v + 1) + c_1u + c_2uv + c_3u^2 = 0$ with $u = \frac{x}{z}$ and $v = \frac{y}{z}$ has a base point $(0, 1)$.

We translate $(0, 1)$ to the origin such that $U_z : c_0v^2 + c_1u + c_2u(v + 1) + c_3u^2 = 0$ and $F(u, v, 1) := F(u, v + 1, 1)$.

We have that $F(uv, v, 1) = c_0v^2 + c_1uv + c_2uv(v+1) + c_3u^2v^2 = v(c_0v + c_1u + c_2u(v+1) + c_3u^2v)$.

It follows that the base point $(0, 1)$ has multiplicity 1.

We continue with $\tilde{U}_{zt} : c_0v + c_1u + c_2u(v+1) + c_3u^2v = 0$ and find base point $(0, 0)$.

We repeat the procedure as above and find that this base point has also multiplicity 1.

There are no other base points.

In order to construct a linear series with a prescribed set of base points, we consider the procedure as above. We want to construct a linear series $\Lambda$ such that the procedure for analyzing base points of $\Lambda$ returns the required result. See chapter 8, section 2 in Lubbes [2011] for the algorithm in pseudo code.

**Example 27. (linear series with prescribed base points)**

We want to create a linear series of degree 2 with base point $(0 : 1 : 1)$ and an infinitely near base point at $(0, 0)$ with respect to the chart $\tilde{U}_{zt}$ as defined above. We want both base points to have multiplicity 1.

We start with the linear series $\langle z^2, yz, y^2, xz, xy, x^2\rangle$.

We consider the linear series $\langle 1, y, y^2, x, xy, x^2\rangle$ for the chart $U_z$.

We move the origin to $(0, 1)$ and obtain $\langle g_0, \ldots, g_5\rangle := \langle 1, y+1, y^2+2y+1, x, xy+x, x^2\rangle$.

We consider the list
$L_0 := \{\ \langle \frac{\partial}{\partial x^a \partial y^b} g_0, \ldots, \frac{\partial}{\partial x^a \partial y^b} g_5\rangle \ \mid\ a+b \leq m-1 \text{ and } a, b \in \mathbf{Z}_{\geq 0}\ \} = \{\langle g_0, \ldots, g_5\rangle\}$ where $m = 1$ is the multiplicity.

We set $\langle h_0, \ldots, h_5\rangle := \langle g_0(xy, y)\backslash y, \ldots, g_5(xy, y)\backslash y\rangle = \langle 0, 1, y+2, x, xy+x, x^2y\rangle$ where $\backslash y$ denotes the polynomial quotient with $y^m$ such that $m = 1$ is the multiplicity. Note that this ensures in Example 26 that the exceptional component $E$ is contained with the right multiplicity.

As above we consider the list for the infinitely base point $(0, 0)$ with multiplicity $n = 1$:
$L_1 := \{\ \langle \frac{\partial}{\partial x^a \partial y^b} h_0, \ldots, \frac{\partial}{\partial x^a \partial y^b} h_5\rangle \ \mid\ a+b \leq n-1 \text{ and } a, b \in \mathbf{Z}_{\geq 0}\ \} = \{\langle h_0, \ldots, h_5\rangle\}$.

The list $L_0 \cup L_1$ defines a matrix of polynomials with rows $(g_i)_i$ and $(h_i)_i$. If we evaluate the polynomials at $(0, 0)$ we obtain the following matrix:

$$M := \begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 & 0 \end{bmatrix}$$

The matrix $M$ expresses the linear conditions on the coefficients of $F(x, y, z) = c_0z^2 + c_1yz + c_2y^2 + c_3xz + c_4xy + c_5x^2 = 0 \subset \mathbf{P}^2$ such that it has a base point at $(0 : 1 : 1)$ and an infinitely near base point as defined before.

The kernel of $M$ is

$$\begin{bmatrix} 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

which corresponds to the linear series $\langle y^2 - 2yz + z^2, xz, xy, x^2\rangle$. Note that this is indeed the linear series which we analyzed in Example 26.

The methods from Example 26 and Example 27 lead to the following algorithms. We have that 'Uz' corresponds to the chart $U_z$ as in the notation above. The chart 'Ct' corresponds to the chart $\tilde{U}_{zt}$.

**Algorithm 28. (base point analysis of linear series in the projective plane)**

```
• function: GetBasePoints( $G$ )
  input: A list $G = (g_i)_i$ such that $g_i \in \mathbf{F}[x, y, z]$ and gcd( $G$ ) $= 1$.
  output: Infinitely near base points of linear series $\Lambda = \langle g_1, \ldots, g_n\rangle$ in the projective plane.
  $L_z$ :=GetBasePointsAffine( $(g_i(x, y, 1))_i$, 0, 'Uz' )
  $L_y$ :=GetBasePointsAffine( $(g_i(x, 1, z))_i$, 0, 'Uy' )
  $L_x$ :=GetBasePointsAffine( $(g_i(1, y, z))_i$, 0, 'Ux' )
  return [ $L_x$, $L_y$, $L_z$ ]
  end function

• function: GetBasePointsAffine( $G$, $depth$, $chart$ )
  input: A list $G = (g_i)_i$ such that $g_i \in \mathbf{F}[u, v]$, gcd( $G$ ) $= 1$, $depth \in \mathbf{Z}_{\geq 0}$ and $chart \in \{$ 'Ux', 'Uy', 'Uz', 'Cs', 'Ct' $\}$.
  output: Infinitely near base points of linear series $\Lambda = \langle g_1, \ldots, g_n\rangle$ in the affine plane.

  % We denote appending elements of an ordered list to another ordered list by $\uplus$
  if $chart$ == 'Ct' then
    $S$ :=ZeroSet( $g_1, \ldots, g_n$, $v$ )
  else if $chart$ == 'Cs' then
    $S$ :=ZeroSet( $g_1, \ldots, g_n$, $u$ )
  else
    $S$ :=ZeroSet( $g_1, \ldots, g_n$)
  end if

  if $S == \emptyset$ then return [ $chart$, $depth$, $\emptyset$, 0, $(g_i)_i$, $\emptyset$, $\emptyset$ ]
  $L := []$
  for $sol \in S$ do
    if ChartOverlap( $sol$, $chart$ ) then
      $L := L \uplus [[\ chart,\ depth,\ sol,\ -1,\ (g_i)_i,\ \emptyset,\ \emptyset\ ]]$
    else
      $(g'_i)_i := (g_i(\ u - sol_u,\ v - sol_v\ ))_i$
      $mul :=$ deg( gcd( $(g'_i(\ uv,\ v\ ))_i$ )

      $L_t$ :=GetBasePointsAffine( $(g'_i(\ uv,\ v\ )v^{-mul})_i$, $depth + 1$, 'Ct' )
      $L_s$ :=GetBasePointsAffine( $(g'_i(\ u,\ vu\ )u^{-mul})_i$, $depth + 1$, 'Cs' )

      $L := L \uplus [[\ chart,\ depth,\ sol,\ mul,\ (g_i)_i,\ L_t,\ L_s\ ]]$
    end if
end for
return $L$
end function
```

- **function:** ChartOverlap( *sol*, *chart* )
  *input:* a solution $sol = (a, b)$ and current chart which is either $Uz, Uy, Uz, Ct$ or $Cs$.
  *output:* True when solution is in one of previous charts, and false otherwise.

```
if $chart ==$ 'Ux' then
  return $False$
else if $chart ==$ 'Uy' then
  return $a \neq 0$
else if $chart ==$ 'Uz' then
  return $a \neq 0$ or $b \neq 0$
else if $chart ==$ 'Ct' then
  return $False$
else if $chart ==$ 'Cs' then
  return $b \neq 0$
end if
end function
```

The algorithm below describes the method used in Example 27 in pseudo code. The 2nd input variable of this algorithm is of type $BPTREE$ as defined below. In Example 27 we have that $degree = 2$ and $[B_z,\ B_y,\ B_x] = [[(0,1), 1, [(0,0), 1], \emptyset], \emptyset, \emptyset]$.

**Algorithm 29. (create linear series in the projective plane with given base points)**

- **function:** GetLinearSeries( *degree*, $[B_z,\ B_y,\ B_x]$ )
  *input:* We need *degree* in $\mathbf{Z}_{\geq 0}$. The lists $B_z, B_y$ and $B_x$ are finite lists of infinitely near base points such that respectively $z = 1, y = 1$ and $x = 1$ and of type $BPTREE$. The recursive type $BPTREE$ is defined as follows: $BPTREE = [[\ sol,\ mult,\ BPTREE,\ BPTREE\ ], \ldots\ ]$ or $BPTREE = \emptyset$ with $sol \in \mathbf{F}^2$ and $mult \in \mathbf{Z}_{>0}$.
  *output:* A linear series $\mathbf{P}(\mathbf{F}\langle g_0, \ldots, g_n\rangle)$ of given degree and with base points defined by $B$ (if such linear series exists).

```
% We denote appending elements of an ordered list to another ordered list by $\uplus$
$(g_i)_i := (\ x^a y^b z^c \ \mid\ a + b + c = degree$ and $a, b, c \in \mathbf{Z}_{\geq 0}\ )$

$L$ :=GetLinearSeriesList( $B_z$, $[(g_i(x, y, 1))_i]$ )
$L$ :=GetLinearSeriesList( $B_y$, $L \uplus [(g_i(x, 1, z))_i]$ )
$(f_{ij})_{ij}$ :=GetLinearSeriesList( $B_x$, $L \uplus [(g_i(1, y, z))_i]$ )

$(k_{ij})_{ij}$ :=Kernel( $(\ f_{ij}(0,0)\ )_{ij}$ )
return $(\ \sum_i k_{ij} g_i\ )_j$
end function
```

- **function:** GetLinearSeriesList( $B$, $L$ )
  *input:* A recursive defined finite list $B$ of type $BPTREE$ and a list of linear series $L = (g_{ij})_{ij}$ with $g_{ij} \in \mathbf{F}[u, v]$.
  *output:* A list $(f_{ij})_{ij}$ of linear series, such that $( f_{ij}(0, 0) )_{ij}$ is a matrix representing the linear conditions on the base points.

```
$(g_i)_i$ :=GetAndRemoveLastElementOfList( $L$ )
for [ $sol$, $mult$, $B_t$, $B_s$] $\in B$ do
  $(f_i)_i := ( g_i( u - sol_u, v - sol_v ) )_i$
  $L := L \uplus [ (\frac{\partial}{\partial u^a \partial v^b} f_i)_i \mid a + b \leq mult - 1 \text{ and } a, b \in \mathbf{Z}_{\geq 0} ]$

  $(a_i)_i$ = (PolynomialQuotient( $h_i(uv, v)$, $v^{mult}$ ))$_i$
  $L$ :=GetLinearSeriesList( $B_t$, $L \uplus [ (a_i)_i ]$ ))

  $(b_i)_i$ = (PolynomialQuotient( $h_i(u, vu)$, $u^{mult}$ ))$_i$
  $L$ :=GetLinearSeriesList( $B_s$, $L \uplus [ (b_i)_i ]$ )
end for
return $L$
end function
```

# 6 Examples of constructing weak Del Pezzo surfaces

For a given geometric C1 label $(L, r)$ in Table 37 we want to construct examples of a weak Del Pezzo surface X in the corresponding Cremona equivalence class. See chapter 7, section 4, remark 98 and remark 99 in Lubbes [2011] for an alternative method for constructing weak Del Pezzo surfaces of degree one and two.

Let $G \xrightarrow{\varphi} R$ be the C1 label elements to root function. We choose a set of points $(p_i)_{i \in [r]}$ in the projective plane which are generic except:

- $p_{i_2}$ is infinitely near to $p_{i_1}$ if and only if $l \in L$ such that $\varphi(l) = Q_{i_1} - Q_{i_2}$,
- $p_{j_t}$ lie on a line if and only if $l \in L$ such that $\varphi(l) = H - Q_{j_1} - Q_{j_2} - Q_{j_3}$,
- $p_{k_t}$ lie on a conic if and only if $l \in L$ such that $\varphi(l) = 2H - Q_{k_1} - Q_{k_2} - Q_{k_3} - Q_{k_4} - Q_{k_5} - Q_{k_6}$, or
- $p_{l_t}$ lie on a cubic with a double point at $p_{l_1}$ if and only if $l \in L$ such that $\varphi(l) = 3H - 2Q_{l_1} - Q_{l_2} - Q_{l_3} - Q_{l_4} - Q_{l_5} - Q_{l_6} - Q_{l_7} - Q_{l_8}$,

for $i_t, j_t, k_t, l_t \in [8]$. Note that for this construction we require that the C1 label $(L, r)$ is geometric.

As in Example 27 we construct a linear series of degree 3 polynomials with base locus $(p_i)_{i \in [r]}$. Let $\mathbf{P}^2 \overset{f}{\dashrightarrow} \mathrm{Y}$ be the associated map of the linear series. We define $\mathrm{X} \xrightarrow{g} \mathbf{P}^2$ to be the blowup of the projective plane at the base locus of the linear series. We define $D \in \mathrm{Pic}\mathrm{X}$ to be the pullback of the hyperplane sections along $g$. We have that $\mathrm{Y} = \varphi_D(\mathrm{X})$

with $D = -K$ being the anticanonical divisor class of X.

$$\begin{tikzcd}
\mathrm{X} \arrow[d, "g"'] \arrow[dr, "\varphi_D"] & \\
\mathbf{P}^2 \arrow[r, dashed, "f"] & \mathrm{Y}
\end{tikzcd}$$

We have that X is a weak Del Pezzo surface of degree $9 - r$ and $D = 3H - Q_1 - \ldots - Q_r$.

**Example 30. (degree four Del Pezzo with two ordinary double points)** Let $(L, r) = (\{\ 1123,\ 45\ \}, 5)$ with Dynkin type $2A_1$ in Table 37 at row index 19.

We choose $p_1 = (0 : 0 : 1)$, $p_2 = (1 : 0 : 1)$ and $p_3 = (2 : 0 : 1)$, such that they lie on the line $y = 0$.

We choose $p_4$ with some random coordinates, say $p_4 = (5 : 7 : 1)$. And finally we take $p_5$ infinitely near to $p_4$ with coordinates $(0, 0)$ in the chart $\tilde{U}'_{zt}$ (see previous section).

Similarly as in Example 27 we construct a linear series of degree 3: $\mathbf{Q}\langle \frac{1}{25}x^2y - \frac{2}{35}xy^2 + yz^2, \frac{7}{25}x^2y - \frac{2}{5}xy^2 + y^2z, \frac{49}{25}x^2y - \frac{14}{5}xy^2 + y^3, \frac{1}{2}x^3 - \frac{1}{2}x^2y - \frac{3}{2}x^2z + \frac{23}{98}xy^2 + xz^2, -\frac{1}{7}xy^2 + xyz \rangle$. This linear series defines a parametrization $\mathbf{P}^2 \overset{f}{\dashrightarrow} \mathrm{Y}$ of a weak degree 4 Del Pezzo surface with $2A_1$ singularities such that $\mathrm{Y} = \varphi_D(\mathrm{X})$. Let $F(\mathrm{X})$ be the Del Pezzo zero-set. Let $\tilde{G} \overset{\varphi}{\to} R$ be the C1 label elements to root function (we use $\tilde{G}$ instead of $G$ since otherwise there is an overlap in notation).

We have that $F_{\geq 0}(\mathrm{X}) = (\varphi(l))_{l \in L} = \{\ H - Q_1 - Q_2 - Q_3,\ Q_4 - Q_5\ \}$ is the indecomposable Del Pezzo zero-set. Let $G(\mathrm{X})$ be the Del Pezzo two-set.

For example $2H - Q_1 - Q_2 - Q_3 - Q_4$ in $G(\mathrm{X})$ and $H - Q_1 - Q_2 - Q_3$ in $F_{\geq 0}(\mathrm{X})$ have intersection product $-1$.

The classes in $G(\mathrm{X})$ which are positive to all classes in $F_{\geq 0}(\mathrm{X})$ is defined by $G_{irr}(\mathrm{X}) = \{\ H - Q_1,\ H - Q_2,\ H - Q_3,\ H - Q_4,\ 2H - Q_1 - Q_2 - Q_4 - Q_5,\ 2H - Q_1 - Q_3 - Q_4 - Q_5,\ 2H - Q_2 - Q_3 - Q_4 - Q_5\ \}$. From Proposition 21 it follows that these classes define linear series without fixed components.

The classes in $G_{irr}(\mathrm{X})$ define the families of conics. So we know that the degree 4 weak Del Pezzo surface Y which we parametrize has 7 families of conics. Similarly we have that the classes in $E_{\geq 0}(\mathrm{X})$ defines the lines of Y, and $F_{\geq 0}(\mathrm{X})$ the isolated singularities.

From Proposition 25 it follows that the geometry determined by the divisor classes is determined up to Cremona equivalence.

**Example 31. (degree one Del Pezzo with four cusps)** Let $(L, r) = (\{\ 1123,\ 1345,\ 1156,\ 1258,\ 1367,\ 1247,\ 1468,\ 1178\ \}, 5)$ with Dynkin type $4A2$ in Table 37 at row index 163.

Suppose we want to plant $n$ trees so that there are 3 trees in each row. The Orchards planting problem is to find a configuration of trees such that the number of rows is maximal.

For $n = 9$ the solution of this problem is called the Pappus configuration, which is depicted in the image below. We find that there are 10 rows, where a row is represented by a line.



The labeled points represent 8 base points in the plane. For example the C1 label element 1123 denotes that a line goes through the base points labeled $1, 2$ and 3. A pair of two lines which do not intersect in the base points form a $A2$ singularity (for example 1123 and 1468). We find three pairs of lines in the Pappus configuration. The fourth pair, 1178 and 1345, are no real lines. It is not possible to choose all points real.

It is well known (see for example Griffiths and Harris [1978], chapter 2, section 4, page 281) that a line through two flexes of a smooth cubic curve passes through a third one. So we can choose the base points on eight of the nine flexes of a cubic plane curve.

The author would like to thank Josef Schicho for pointing out the Pappus configuration.

# 7 Algorithm: real structures of weak Del Pezzo surfaces

In this section we present an algorithm for classifying the real structures of weak Del Pezzo surfaces up to conjugacy. The classification of real structures of Del Pezzo surfaces is well known and can be found in Wall [1987]. See also Manin [1966] and Silhol [1989].

The correctness of our algorithm is based on the fact that the conjugacy classes of real structures are uniquely determined by the Weyl equivalence classes of root subsystems. We can use the classification of C1 root subsystems.

As an application we consider conic families on real weak Del Pezzo surfaces of degree 4 in Example 36.

**Definition 32. (real structure for weak Del Pezzo surfaces)** Let $Gal(\mathbf{C}|\mathbf{R}) = \langle\sigma\rangle$ be the Galois group. We define a *real weak Del Pezzo surface* to be a pair $(\mathrm{X}, \sigma)$ where X is a complex weak Del Pezzo surface, and $\sigma(\mathrm{X}) = \mathrm{X}$. Let $A(\mathrm{X}) = (\ \mathrm{Pic}(\mathrm{X}),\ K,\ \cdot,\ h\ )$ be the enhanced Picard group. We define a *real structure* of $(\mathrm{X}, \sigma)$ to be the automorphism of the enhanced Picard group induced by $\sigma$. By abuse of notation we will denote the real structure also by $\sigma$. We have that $\sigma(-K) = -K$ where $-K$ is the anticanonical divisor class of X.

**Algorithm 33. (real structures of weak Del Pezzo surfaces)**

*% With square brackets we denote ordered and indexed lists. We denote appending elements of an ordered list to another ordered list by* $\uplus$. *We represent a C1 label* $(L, r)$ *as a set of column vectors of length* $r+1$ *with respect to the standard Del Pezzo basis. We will represent the equivalence class of a real structure by a matrix acting on column vectors. We represent C1 labels also by using the C1 label elements to root function, defined in Definition 8. Thus for example* 1123 *represents the column vector* $(1, -1, -1, -1, 0, \dots, 0)$.

- **function:** getDPRealStructures()
  *output:* Returns a list of unique representatives of equivalence classes of all real structures for weak Del Pezzo surfaces. The list consists of entries $(L, M)$ where $M$ is a matrix representing a real structure with respect to the standard Del Pezzo basis and $L$ is a C1 label, which represents the corresponding root subsystem.

```
$C1$ :=getGeometricC1RootSubsystemsClassification() (see Algorithm 11)
$T = []$
for $(L, rank) \in C1$ do
  if $L \subset \{1123, 12, 23, 34, 45, 56, 67, 78\}$ then
    $V$ :=extendToBasis($L$)
    $D$ :=getRealEigenValueMatrix( rank($V$), length($L$) )
    $M := V \cdot D \cdot V^{-1}$
    if isIntegralMatrix($M$) then
      $T \uplus [\ (L, M)\ ]$
    end if
  end if
end for
return $T$
end function
```

- **function:** getRealEigenValueMatrix($rank, n$)
  *input:* Integers $rank$ and $n$.
  *output:* A square diagonal matrix of given $rank$ where the first $n$ diagonal entries are $-1$, and 1 otherwise.
  **end function**

- **function:** extendToBasis($L$)
  *input:* A matrix $L$, with linear independent column vectors corresponding to a C1 label of types { 1123, 12, 23, 34, 45, 56, 67, 78 }.
  *output:* A matrix forming a basis consisting of the column vectors $L$ appended by column vectors which are orthogonal to $L$ with respect to intersection product as in Definition 6.

```
$A$ :=sortColumnsLexicographic($L$)
$has1123 := 0$
if $A_{0,0} == 1$ then
  removeFirstColumnFrom($A$)
  $has1123 := 1$
end if
$E := []$
for $zeroRow \in$ zeroRowVectorsOf($A$) do
  if indexOf($zeroRow$)!=$has1123 - 1$ then
    $v$ :=canonicalBasisColumnVector( indexOf($zeroRow$) )
    $v_0 := -has1123\cdot$intersectionProduct($v, 1123$)
    $E := E \uplus [v]$
  end if
end for
while not isSquareMatrix($L \cup E$) do
  $C$ := [ getAndRemoveFirstColumnFrom($A$) ]
  while not isOrthogonal( lastColumnOf($C$), firstColumnOf($A$) ) do
    $C := C \uplus$ [ getAndRemoveFirstColumnFrom($A$) ]
  end while
  $a$ :=firstNonZeroRowIndex( firstElementOf($C$) )
  $b$ :=lastNonZeroRowIndex( lastElementOf($C$) )
  $v$ :=zeroColumnVector()
  $v_a := 1$; $v_{a+1} := 1$; $\ldots$; $v_b := 1$
  $v_0 := -has1123\cdot$intersectionProduct($v, 1123$)
  $E := E \uplus [v]$
end while
return matrix($L \cup E$)
end function
```

**Example 34. (real structures of weak Del Pezzo surfaces)** Let $(L, r) = (\{\ 1123,\ 12,\ 23,\ 45\ \}, 5)$ be a C1 label of type $A_2 + 2A_1$ (index 26 in Table 37). We trace the getDPRealStructures() function in Algorithm 33 line by line for this C1 label.

Using vector notation with the standard Del Pezzo basis as in Definition 6 we represent $L$ as $[(1, -1, -1, -1, 0, 0), (0, 1, -1, 0, 0, 0), (0, 0, 1, -1, 0, 0), (0, 0, 0, 0, 1, -1)]$.

We have that $L \subset \{1123, 12, 23, 34, 45, 56, 67, 78\}$.

We have that

$$V = \begin{bmatrix} 1 & 0 & 0 & 0 & -3 & 0 \\ -1 & 1 & 0 & 0 & 1 & 0 \\ -1 & -1 & 1 & 0 & 1 & 0 \\ -1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 1 \end{bmatrix}, D = \begin{bmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, M = \begin{bmatrix} 2 & 1 & 1 & 1 & 0 & 0 \\ -1 & -\frac{4}{3} & -\frac{1}{3} & -\frac{1}{3} & 0 & 0 \\ -1 & -\frac{1}{3} & -\frac{4}{3} & -\frac{1}{3} & 0 & 0 \\ -1 & -\frac{1}{3} & -\frac{1}{3} & -\frac{4}{3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}.$$

We find that $M$ is not an integral matrix and thus not a valid real structure.

**Theorem 35. (real structures of weak Del Pezzo surfaces)**

**a)** Algorithm 33 is correct.

**Proof:**

*Claim 1:* We have that the extendToBasis$(L)$ function is correct.
Left to the reader.

Let $(\mathrm{X}, \sigma)$ be a real weak Del Pezzo surface. Let $A(\mathrm{X}) = (\ \mathrm{Pic}(\mathrm{X}),\ K,\ \cdot,\ h\ )$ be the enhanced Picard group of X. Let $(\tilde{M}, (\cdot,\cdot), <, R)$ be a C1 kit such that $A(\mathrm{X}) \cong (\tilde{M}, (-3, 1, \ldots, 1), (\cdot,\cdot))$. Let $D$, $M$ and $V$ be the matrices as in Algorithm 33.

*Claim 2:* We have that **a)**.
A real structure of a weak Del Pezzo surface induces an involution $\sigma$ on $W$ such that $W = W^+ \oplus W^-$ where $W^+ = \{\ w \in W \ \mid\ \sigma(w) = w\ \}$ and $W^- = \{\ w \in W \ \mid\ \sigma(w) = -w\ \}$. From $\sigma(K) = K$ and $CK = \sigma(C)\sigma(K) = -CK$ for $C \in W^-$ it follows that $W^-$ is orthogonal to $K$. From Proposition 7 it follows that $W^-$ is generated by a subset of a basis for the root system $R$. It follows that $W^-$ is generated by a basis of a root subsystem of $R$ such that the basis is a subset of $\{1123, 12, 23, 34, 45, 56, 67, 78\}$. Up to conjugacy we only have to consider the root subsystems of the C1 classification. The eigenvectors of the eigenvalue $-1$ are determined by a C1 label. The hyperplanes orthogonal to the roots in $W^-$ are point wise fixed. The eigenvectors of the eigenvalue 1 are linear independent vectors which are orthogonal to eigenspace of $-1$. It follows that $MV = VD$ and thus $M = VDV^{-1}$. Since $M$ should induce a $\mathbf{Z}$-module automorphism we need that the matrix $M$ is integral. ☕

The output of Algorithm 33 is in Table 38. Note that we observe from Table 38 that $(rank, f0, f1)$ is a strong invariant for the real structure. As an application of the classification of real structures in explicit coordinates we consider weak Del Pezzo surfaces of degree 4.

**Example 36. (real structures of weak Del Pezzo surface of degree four)** Let $(\mathrm{X}, \sigma)$ be a real weak Del Pezzo surface of degree 4. Let $C$ be the set of C1 labels of root subsystems of $D_5$ (indices $16, \ldots, 31$ in Table 37). Let $R$ be the set of conjugacy classes of real structures of X (indices $10, \ldots, 15$ in Table 38). Let $B$ be the set of subsets of the positive roots in the Del Pezzo zero-set of X. Let $B(c) = \{\ b \in B \ \mid\ \mathrm{isomorphicBases}(c, b) == True\ \}$ for $c \in C$ (see Algorithm 11). Let $B(c, r) = \{\ b \in B(c) \ \mid\ r(b) = b\ \}$ for $r \in R$ (note that we need that the basis of a root subsystem is compatible with the real structure). Let $G(c, r, b)$ be the set of classes in the irreducible Del Pezzo two-set which are fixed by the real structure $r \in R$ and where the real irreducible Del Pezzo zero-set is $b \in B(c, r)$ The following table denotes the possible cardinalities of $G(c, r, b)$ by fixing a representative $r$ for a conjugacy class of a real structure in $R$, and considering all bases in $b \in B(c)$ of the same Dynkin type as $c$. The 'index' column denotes the index of $c$ in Table 37 and the 'type' column, the Dynkin type of $c$. The 'index' row denote the index $i$ for the real structure $t$ in Table 38 and the 'type' row, the Dynkin type of $t$.

| index | | 10 | 11 | 12 | 13 | 14 | 15 |
|---|---|---|---|---|---|---|---|
| | type | $A0$ | $A1$ | $2A1$ | $2A1$ | $3A1$ | $D4$ |
| 16 | $A0$ | 10 | 6 | 2 | 6 | 2 | 2 |
| 17 | $A1$ | 8 | 4, 6 | 2 | 4 | 2 | |
| 18 | $2A1$ | 6 | 4 | 2 | | | |
| 19 | $2A1$ | 7 | 3 | | 3, 5 | 1 | 1 |
| 20 | $A2$ | 6 | 2 | | 2 | 2 | |
| 21 | $3A1$ | 5 | 3 | | 3 | 1 | |
| 22 | $A2 + A1$ | 4 | 2 | | | | |
| 23 | $A3$ | 4 | | 2 | | | |
| 24 | $A3$ | 5 | 1 | | 1, 3 | | |
| 25 | $4A1$ | 4 | | 2 | 2, 4 | | |
| 26 | $A2 + 2A1$ | 3 | | | | 1 | |
| 27 | $A3 + A1$ | 3 | 1 | | | | |
| 28 | $A4$ | 2 | | | | | |
| 29 | $D4$ | 3 | | | 1 | | |
| 30 | $A3 + 2A1$ | 2 | | | 2 | | |
| 31 | $D5$ | 1 | | | | | |

The geometric interpretation is that each element of $G(c, r, b)$ is the class of a one dimensional family of conics on X. For example a real torus in three space has 4 complex $A_1$ singularities and 4 families of circles. The singularities and the real structure of the torus are defined by entry $(row, column) = (25, 10)$ in the above table. We see from the table that the number of families of conics on X is not uniquely defined by only the conjugacy classes of $c$ and $t$. For example if $(row, column) = (17, 11)$ then there exists $b = [12], b' = [1123] \in B(c)$ such that $\#G(c, r, b) = 4$ and $\#G(c, r, b') = 6$. We have that $r : (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5) = \alpha_0 H + \sum \alpha_i Q_i \mapsto (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_4)$, $G(c, r, b) = \{\ H - Q_1,$ $H - Q_3,\ 2H - Q_1 - Q_2 - Q_4 - Q_5,\ 2H - Q_1 - Q_3 - Q_4 - Q_5\ \}$ and $G(c, r, b') = \{\ H - Q_1,$ $H - Q_2,\ H - Q_3,\ 2H - Q_1 - Q_2 - Q_4 - Q_5,\ 2H - Q_1 - Q_3 - Q_4 - Q_5,\ 2H - Q_2 - Q_3 - Q_4 - Q_5$ $\}$. The obvious algorithm computing the above table does not terminate in reasonable time for degree one and degree two weak Del Pezzo surfaces, since $B$ is too large. The roots in the eigenspace of $-1$ cannot be effective. It follows that we only have to consider bases $b \in B$ which are disjoint to the eigenspace of $-1$.

## 8 Acknowledgements

It is my pleasure to acknowledge that the many computations with Josef Schicho is a major contribution to this paper. Also I would like to thank Michael Harrison for useful discussions concerning root subsystems.

The algorithms were implemented using the computer algebra system Sage (Stein et al. [2012]) with additional interfacing to Magma (Bosma et al. [1997]).

This research was supported by the Austrian Science Fund (FWF): project P21461.

# A Table of geometric C1 labels

**Table 37. (geometric bases of C1 root subsystems up to Weyl equivalence)**

- the ‘index’ column assigns an index to each row for future reference,
- the ‘rank’ column defines the rank $r$ of the C1 root system,
- the ‘basis’ column denotes the basis $L$ such that $(L, r)$ is a geometric C1 label,
- the ‘type’ column denotes the Dynkin type of the C1 label $(L, r)$.

| index | rank | basis | type |
|---|---|---|---|
| 1 | 2 | [] | $A0$ |
| 2 | 2 | [12] | $A1$ |
| 3 | 3 | [] | $A0$ |
| 4 | 3 | [23] | $A1$ |
| 5 | 3 | [1123] | $A1$ |
| 6 | 3 | [1123, 23] | $2A1$ |
| 7 | 3 | [12, 23] | $A2$ |
| 8 | 3 | [1123, 12, 23] | $A2 + A1$ |
| 9 | 4 | [] | $A0$ |
| 10 | 4 | [34] | $A1$ |
| 11 | 4 | [12, 34] | $2A1$ |
| 12 | 4 | [23, 34] | $A2$ |
| 13 | 4 | [1123, 12, 34] | $A2 + A1$ |
| 14 | 4 | [12, 23, 34] | $A3$ |
| 15 | 4 | [1123, 12, 23, 34] | $A4$ |
| 16 | 5 | [] | $A0$ |
| 17 | 5 | [45] | $A1$ |
| 18 | 5 | [23, 45] | $2A1$ |
| 19 | 5 | [1123, 45] | $2A1$ |
| 20 | 5 | [34, 45] | $A2$ |
| 21 | 5 | [1123, 23, 45] | $3A1$ |
| 22 | 5 | [12, 34, 45] | $A2 + A1$ |
| 23 | 5 | [23, 34, 45] | $A3$ |
| 24 | 5 | [1123, 34, 45] | $A3$ |
| 25 | 5 | [1145, 1123, 23, 45] | $4A1$ |
| 26 | 5 | [1123, 12, 23, 45] | $A2 + 2A1$ |
| 27 | 5 | [1123, 12, 34, 45] | $A3 + A1$ |
| 28 | 5 | [12, 23, 34, 45] | $A4$ |
| 29 | 5 | [1123, 23, 34, 45] | $D4$ |
| 30 | 5 | [1145, 1123, 12, 23, 45] | $A3 + 2A1$ |
| 31 | 5 | [1123, 12, 23, 34, 45] | $D5$ |
| 32 | 6 | [] | $A0$ |
| 33 | 6 | [56] | $A1$ |
| 34 | 6 | [34, 56] | $2A1$ |
| 35 | 6 | [45, 56] | $A2$ |
| 36 | 6 | [12, 34, 56] | $3A1$ |
| 37 | 6 | [23, 45, 56] | $A2 + A1$ |
| 38 | 6 | [34, 45, 56] | $A3$ |
| 39 | 6 | [1145, 1123, 23, 45] | $4A1$ |
| 40 | 6 | [1123, 23, 45, 56] | $A2 + 2A1$ |
| 41 | 6 | [12, 23, 45, 56] | $2A2$ |
| 42 | 6 | [12, 34, 45, 56] | $A3 + A1$ |
| 43 | 6 | [23, 34, 45, 56] | $A4$ |
| 44 | 6 | [1123, 23, 34, 45] | $D4$ |
| 45 | 6 | [1123, 12, 23, 45, 56] | $2A2 + A1$ |
| 46 | 6 | [1145, 1123, 23, 45, 56] | $A3 + 2A1$ |
| 47 | 6 | [1123, 12, 34, 45, 56] | $A4 + A1$ |
| 48 | 6 | [12, 23, 34, 45, 56] | $A5$ |
| 49 | 6 | [1123, 23, 34, 45, 56] | $D5$ |
| 50 | 6 | [1456, 1123, 12, 23, 45, 56] | $3A2$ |
| 51 | 6 | [1145, 1123, 12, 23, 45, 56] | $A5 + A1$ |
| 52 | 6 | [1123, 12, 23, 34, 45, 56] | $E6$ |
| 53 | 7 | [] | $A0$ |
| 54 | 7 | [67] | $A1$ |
| 55 | 7 | [45, 67] | $2A1$ |
| 56 | 7 | [56, 67] | $A2$ |
| 57 | 7 | [23, 45, 67] | $3A1$ |
| 58 | 7 | [1123, 45, 67] | $3A1$ |
| 59 | 7 | [34, 56, 67] | $A2 + A1$ |
| 60 | 7 | [45, 56, 67] | $A3$ |
| 61 | 7 | [1123, 23, 45, 67] | $4A1$ |
| 62 | 7 | [1145, 1123, 23, 45] | $4A1$ |

| | | | |
|---|---|---|---|
| 63 | 7 | [12, 34, 56, 67] | $A2 + 2A1$ |
| 64 | 7 | [23, 34, 56, 67] | $2A2$ |
| 65 | 7 | [23, 45, 56, 67] | $A3 + A1$ |
| 66 | 7 | [1123, 45, 56, 67] | $A3 + A1$ |
| 67 | 7 | [34, 45, 56, 67] | $A4$ |
| 68 | 7 | [1123, 23, 34, 45] | $D4$ |
| 69 | 7 | [1145, 1123, 23, 45, 67] | $5A1$ |
| 70 | 7 | [1123, 12, 23, 45, 67] | $A2 + 3A1$ |
| 71 | 7 | [1123, 12, 34, 56, 67] | $2A2 + A1$ |
| 72 | 7 | [1123, 23, 45, 56, 67] | $A3 + 2A1$ |
| 73 | 7 | [1145, 1123, 23, 45, 56] | $A3 + 2A1$ |
| 74 | 7 | [12, 23, 45, 56, 67] | $A3 + A2$ |
| 75 | 7 | [12, 34, 45, 56, 67] | $A4 + A1$ |
| 76 | 7 | [23, 34, 45, 56, 67] | $A5$ |
| 77 | 7 | [1123, 34, 45, 56, 67] | $A5$ |
| 78 | 7 | [1123, 23, 34, 45, 67] | $D4 + A1$ |
| 79 | 7 | [1123, 23, 34, 45, 56] | $D5$ |
| 80 | 7 | [1567, 1347, 1127, 12, 34, 56] | $6A1$ |
| 81 | 7 | [1456, 1123, 12, 23, 45, 56] | $3A2$ |
| 82 | 7 | [1145, 1123, 12, 23, 45, 67] | $A3 + 3A1$ |
| 83 | 7 | [1123, 12, 23, 45, 56, 67] | $A3 + A2 + A1$ |
| 84 | 7 | [278, 12, 23, 34, 56, 67] | $2A3$ |
| 85 | 7 | [1123, 12, 23, 34, 56, 67] | $A4 + A2$ |
| 86 | 7 | [1123, 12, 34, 45, 56, 67] | $A5 + A1$ |
| 87 | 7 | [1145, 1123, 12, 23, 45, 56] | $A5 + A1$ |
| 88 | 7 | [12, 23, 34, 45, 56, 67] | $A6$ |
| 89 | 7 | [1145, 1123, 23, 45, 56, 67] | $D4 + 2A1$ |
| 90 | 7 | [1123, 12, 23, 34, 45, 67] | $D5 + A1$ |
| 91 | 7 | [1123, 23, 34, 45, 56, 67] | $D6$ |
| 92 | 7 | [1123, 12, 23, 34, 45, 56] | $E6$ |
| 93 | 7 | [278, 1567, 1347, 1127, 12, 34, 56] | $7A1$ |
| 94 | 7 | [278, 1567, 12, 23, 34, 56, 67] | $2A3 + A1$ |
| 95 | 7 | [1456, 1123, 12, 23, 45, 56, 67] | $A5 + A2$ |
| 96 | 7 | [278, 12, 23, 34, 45, 56, 67] | $A7$ |
| 97 | 7 | [278, 1347, 1127, 12, 34, 45, 56] | $D4 + 3A1$ |
| 98 | 7 | [1145, 1123, 12, 23, 45, 56, 67] | $D6 + A1$ |
| 99 | 7 | [1123, 12, 23, 34, 45, 56, 67] | $E7$ |

| | | | |
|---|---|---|---|
| 100 | 8 | [] | $A0$ |
| 101 | 8 | [78] | $A1$ |
| 102 | 8 | [56, 78] | $2A1$ |
| 103 | 8 | [67, 78] | $A2$ |
| 104 | 8 | [34, 56, 78] | $3A1$ |
| 105 | 8 | [45, 67, 78] | $A2 + A1$ |
| 106 | 8 | [56, 67, 78] | $A3$ |
| 107 | 8 | [12, 34, 56, 78] | $4A1$ |
| 108 | 8 | [1145, 1123, 23, 45] | $4A1$ |
| 109 | 8 | [23, 45, 67, 78] | $A2 + 2A1$ |
| 110 | 8 | [34, 45, 67, 78] | $2A2$ |
| 111 | 8 | [34, 56, 67, 78] | $A3 + A1$ |
| 112 | 8 | [45, 56, 67, 78] | $A4$ |
| 113 | 8 | [1123, 23, 34, 45] | $D4$ |
| 114 | 8 | [1145, 1123, 23, 45, 78] | $5A1$ |
| 115 | 8 | [1123, 23, 45, 67, 78] | $A2 + 3A1$ |
| 116 | 8 | [12, 34, 45, 67, 78] | $2A2 + A1$ |
| 117 | 8 | [12, 34, 56, 67, 78] | $A3 + 2A1$ |
| 118 | 8 | [1145, 1123, 23, 45, 56] | $A3 + 2A1$ |
| 119 | 8 | [23, 34, 56, 67, 78] | $A3 + A2$ |
| 120 | 8 | [23, 45, 56, 67, 78] | $A4 + A1$ |
| 121 | 8 | [34, 45, 56, 67, 78] | $A5$ |
| 122 | 8 | [1123, 23, 34, 45, 78] | $D4 + A1$ |
| 123 | 8 | [1123, 23, 34, 45, 56] | $D5$ |
| 124 | 8 | [1567, 1347, 1127, 12, 34, 56] | $6A1$ |
| 125 | 8 | [1145, 1123, 23, 45, 67, 78] | $A2 + 4A1$ |
| 126 | 8 | [1123, 12, 23, 45, 67, 78] | $2A2 + 2A1$ |
| 127 | 8 | [1456, 1123, 12, 23, 45, 56] | $3A2$ |
| 128 | 8 | [1145, 1123, 23, 45, 56, 78] | $A3 + 3A1$ |
| 129 | 8 | [1123, 12, 34, 56, 67, 78] | $A3 + A2 + A1$ |
| 130 | 8 | [12, 23, 34, 56, 67, 78] | $2A3$ |
| 131 | 8 | [1567, 1145, 1127, 12, 56, 78] | $2A3$ |
| 132 | 8 | [1123, 23, 45, 56, 67, 78] | $A4 + 2A1$ |
| 133 | 8 | [12, 23, 45, 56, 67, 78] | $A4 + A2$ |
| 134 | 8 | [12, 34, 45, 56, 67, 78] | $A5 + A1$ |
| 135 | 8 | [1145, 1123, 12, 23, 45, 56] | $A5 + A1$ |
| 136 | 8 | [23, 34, 45, 56, 67, 78] | $A6$ |
| 137 | 8 | [1145, 1123, 23, 45, 56, 67] | $D4 + 2A1$ |
| 138 | 8 | [1123, 23, 34, 45, 67, 78] | $D4 + A2$ |
| 139 | 8 | [1123, 23, 34, 45, 56, 78] | $D5 + A1$ |
| 140 | 8 | [1123, 23, 34, 45, 56, 67] | $D6$ |
| 141 | 8 | [1123, 12, 23, 34, 45, 56] | $E6$ |
| 142 | 8 | [278, 1567, 1347, 1127, 12, 34, 56] | $7A1$ |
| 143 | 8 | [1456, 1123, 12, 23, 45, 56, 78] | $3A2 + A1$ |

| | | | |
|---|---|---|---|
| 144 | 8 | [278, 1347, 1127, 12, 34, 56, 78] | $A3 + 4A1$ |
| 145 | 8 | [1145, 1123, 12, 23, 45, 67, 78] | $A3 + A2 + 2A1$ |
| 146 | 8 | [1678, 1456, 1347, 1123, 12, 45, 78] | $2A3 + A1$ |
| 147 | 8 | [1123, 12, 23, 45, 56, 67, 78] | $A4 + A2 + A1$ |
| 148 | 8 | [1123, 12, 23, 34, 56, 67, 78] | $A4 + A3$ |
| 149 | 8 | [1145, 1123, 12, 23, 45, 56, 78] | $A5 + 2A1$ |
| 150 | 8 | [1456, 1123, 12, 23, 45, 56, 67] | $A5 + A2$ |
| 151 | 8 | [1123, 12, 34, 45, 56, 67, 78] | $A6 + A1$ |
| 152 | 8 | [12, 23, 34, 45, 56, 67, 78] | $A7$ |
| 153 | 8 | [1567, 1145, 1127, 12, 23, 56, 78] | $A7$ |
| 154 | 8 | [1567, 1347, 1127, 12, 34, 56, 78] | $D4 + 3A1$ |
| 155 | 8 | [1567, 1347, 1145, 1127, 12, 56, 78] | $D4 + A3$ |
| 156 | 8 | [1145, 1123, 23, 45, 56, 67, 78] | $D5 + 2A1$ |
| 157 | 8 | [1123, 12, 23, 34, 45, 67, 78] | $D5 + A2$ |
| 158 | 8 | [1145, 1123, 12, 23, 45, 56, 67] | $D6 + A1$ |
| 159 | 8 | [1123, 23, 34, 45, 56, 67, 78] | $D7$ |
| 160 | 8 | [1123, 12, 23, 34, 45, 56, 78] | $E6 + A1$ |
| 161 | 8 | [1123, 12, 23, 34, 45, 56, 67] | $E7$ |
| 162 | 8 | [308, 278, 1567, 1347, 1127, 12, 34, 56] | $8A1$ |
| 163 | 8 | [1123, 1345, 1156, 1258, 1367, 1247, 1468, 1178] | $4A2$ |
| 164 | 8 | [308, 278, 1567, 12, 23, 34, 56, 67] | $2A3 + 2A1$ |
| 165 | 8 | [278, 1678, 12, 23, 34, 45, 67, 78] | $2A4$ |
| 166 | 8 | [1678, 1145, 1123, 12, 23, 45, 67, 78] | $A5 + A2 + A1$ |
| 167 | 8 | [234, 1145, 1123, 12, 23, 56, 67, 78] | $A7 + A1$ |
| 168 | 8 | [1567, 1123, 12, 23, 34, 56, 67, 78] | $A8$ |
| 169 | 8 | [278, 1567, 1347, 1127, 12, 34, 56, 78] | $D4 + 4A1$ |
| 170 | 8 | [234, 278, 12, 23, 34, 56, 67, 78] | $2D4$ |
| 171 | 8 | [278, 1567, 12, 23, 34, 56, 67, 78] | $D5 + A3$ |
| 172 | 8 | [278, 1347, 1127, 12, 34, 45, 56, 78] | $D6 + 2A1$ |
| 173 | 8 | [278, 12, 23, 34, 45, 56, 67, 78] | $D8$ |
| 174 | 8 | [1456, 1123, 12, 23, 45, 56, 67, 78] | $E6 + A2$ |
| 175 | 8 | [1145, 1123, 12, 23, 45, 56, 67, 78] | $E7 + A1$ |
| 176 | 8 | [1123, 12, 23, 34, 45, 56, 67, 78] | $E8$ |

# B Real structures of weak Del Pezzo surfaces

The following table represents the output of Algorithm 33.

**Table 38. (real structures of weak Del Pezzo surfaces)**

- the 'index' column denotes the row index for future reference,
- the 'rank' column denotes that the root subsystem of the real structure is contained in a root system of given rank,
- the 'C1' column denotes the index in Table 37 corresponding to the root subsystem of the real structure,
- the 'type' column denotes the type of the root root subsystem of the real structure,
- the 'f0' columns denote the cardinality of the real Del Pezzo zero-set (thus classes in the Del Pezzo zero-set which are fixed by the real structure),
- the 'f1' columns denote the cardinality of the real Del Pezzo one-set, and
- the 'f2' columns denote the cardinality of the real Del Pezzo two-set.

| index | rank | C1 | type | f0 | f1 | f2 |
|---|---|---|---|---|---|---|
| 1 | 2 | 1 | $A0$ | 2 | 3 | 2 |
| 2 | 2 | 2 | $A1$ | 0 | 1 | 0 |
| 3 | 3 | 3 | $A0$ | 8 | 6 | 3 |
| 4 | 3 | 4 | $A1$ | 2 | 2 | 1 |
| 5 | 3 | 5 | $A1$ | 6 | 0 | 3 |
| 6 | 3 | 6 | $2A1$ | 0 | 0 | 1 |
| 7 | 4 | 9 | $A0$ | 20 | 10 | 5 |
| 8 | 4 | 10 | $A1$ | 6 | 4 | 3 |
| 9 | 4 | 11 | $2A1$ | 0 | 2 | 1 |
| 10 | 5 | 16 | $A0$ | 40 | 16 | 10 |
| 11 | 5 | 17 | $A1$ | 14 | 8 | 6 |
| 12 | 5 | 18 | $2A1$ | 4 | 4 | 2 |
| 13 | 5 | 19 | $2A1$ | 12 | 0 | 6 |
| 14 | 5 | 21 | $3A1$ | 2 | 0 | 2 |
| 15 | 5 | 29 | $D4$ | 0 | 0 | 2 |
| 16 | 6 | 32 | $A0$ | 72 | 27 | 27 |
| 17 | 6 | 33 | $A1$ | 30 | 15 | 15 |
| 18 | 6 | 34 | $2A1$ | 12 | 7 | 7 |
| 19 | 6 | 36 | $3A1$ | 2 | 3 | 3 |
| 20 | 6 | 44 | $D4$ | 0 | 3 | 3 |
| 21 | 7 | 53 | $A0$ | 126 | 56 | 126 |
| 22 | 7 | 54 | $A1$ | 60 | 32 | 60 |
| 23 | 7 | 55 | $2A1$ | 26 | 16 | 26 |
| 24 | 7 | 57 | $3A1$ | 8 | 8 | 8 |
| 25 | 7 | 58 | $3A1$ | 24 | 0 | 24 |
| 26 | 7 | 61 | $4A1$ | 6 | 0 | 6 |
| 27 | 7 | 68 | $D4$ | 6 | 8 | 6 |
| 28 | 7 | 78 | $D4 + A1$ | 4 | 0 | 4 |
| 29 | 7 | 91 | $D6$ | 2 | 0 | 2 |
| 30 | 7 | 99 | $E7$ | 0 | 0 | 0 |
| 31 | 8 | 100 | $A0$ | 240 | 240 | 2160 |
| 32 | 8 | 101 | $A1$ | 126 | 126 | 756 |
| 33 | 8 | 102 | $2A1$ | 60 | 60 | 252 |
| 34 | 8 | 104 | $3A1$ | 26 | 26 | 72 |
| 35 | 8 | 107 | $4A1$ | 8 | 8 | 24 |
| 36 | 8 | 113 | $D4$ | 24 | 24 | 24 |
| 37 | 8 | 122 | $D4 + A1$ | 6 | 6 | 12 |
| 38 | 8 | 140 | $D6$ | 4 | 4 | 4 |
| 39 | 8 | 161 | $E7$ | 2 | 2 | 0 |
| 40 | 8 | 176 | $E8$ | 0 | 0 | 0 |

Below is an explicit coordinate description of the real structure for each index in the above table:

$$\sigma : A(\mathrm{X}) \to A(\mathrm{X}), \quad (H, Q_1, \ldots, Q_r) \mapsto (D_0, \ldots, D_r)$$

where $A(\mathrm{X})$ is the enhanced Picard group of a real weak Del Pezzo surface $(\mathrm{X}, \sigma)$. The coordinates are with respect to the standard Del Pezzo basis $B(r)$. Below we denote $(D_0, \ldots, D_r)$ for each index.$\bullet$ $index = 1$, $( H , Q_1 , Q_2 )$ $\bullet$ $index = 2$, $( H , Q_2 , Q_1 )$ $\bullet$ $index = 3$, $( H , Q_1 , Q_2 , Q_3 )$ $\bullet$ $index = 4$, $( H , Q_1 , Q_3 , Q_2 )$ $\bullet$ $index = 5$, $( 2H - Q_1 - Q_2 - Q_3 , H - Q_2 - Q_3 , H - Q_1 - Q_3 , H - Q_1 - Q_2 )$ $\bullet$ $index = 6$, $( 2H - Q_1 - Q_2 - Q_3 , H - Q_2 - Q_3 , H - Q_1 - Q_2 , H - Q_1 - Q_3 )$ $\bullet$ $index = 7$, $( H , Q_1 , Q_2 , Q_3 , Q_4 )$ $\bullet$ $index = 8$, $( H , Q_1 , Q_2 , Q_4 , Q_3 )$ $\bullet$ $index = 9$, $( H , Q_2 , Q_1 , Q_4 , Q_3 )$ $\bullet$ $index = 10$, $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 )$ $\bullet$ $index = 11$, $( H , Q_1 , Q_2 , Q_3 , Q_5 , Q_4 )$ $\bullet$ $index = 12$, $( H , Q_1 , Q_3 , Q_2 , Q_5 , Q_4 )$ $\bullet$ $index = 13$, $( 2H - Q_1 - Q_2 - Q_3 , H - Q_2 - Q_3 , H - Q_1 - Q_3 , H - Q_1 - Q_2 , Q_5 , Q_4 )$ $\bullet$ $index = 14$, $( 2H - Q_1 - Q_2 - Q_3 , H - Q_2 - Q_3 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , Q_5 , Q_4 )$ $\bullet$ $index = 15$, $( 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , 2H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 )$ $\bullet$ $index = 16$, $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 , Q_6 )$ $\bullet$ $index = 17$, $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_6 , Q_5 )$ $\bullet$ $index = 18$, $( H , Q_1 , Q_2 , Q_4 , Q_3 , Q_6 , Q_5 )$ $\bullet$ $index = 19$, $( H , Q_2 , Q_1 , Q_4 , Q_3 , Q_6 , Q_5 )$ $\bullet$ $index = 20$, $( 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , 2H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , Q_6 )$ $\bullet$ $index = 21$, $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 , Q_6 , Q_7 )$ $\bullet$ $index = 22$, $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 , Q_7 , Q_6 )$ $\bullet$ $index = 23$, $( H , Q_1 , Q_2 , Q_3 , Q_5 , Q_4 , Q_7 , Q_6 )$ $\bullet$ $index = 24$, $( H , Q_1 , Q_3 , Q_2 , Q_5 , Q_4 , Q_7 , Q_6 )$ $\bullet$ $index = 25$, $( 2H - Q_1 - Q_2 - Q_3 , H - Q_2 - Q_3 , H - Q_1 - Q_3 , H - Q_1 - Q_2 , Q_5 , Q_4 , Q_7 , Q_6 )$ $\bullet$ $index = 26$, $( 2H - Q_1 - Q_2 - Q_3 , H - Q_2 - Q_3 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , Q_5 , Q_4 , Q_7 , Q_6 )$ $\bullet$ $index = 27$, $( 3H - 2Q_1 - Q_2 - Q_3 - Q_4 -$

$Q_5 , 2H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , Q_6 , Q_7 )$ $\bullet$ $\mathit{index} = 28,$ $( 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , 2H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , Q_7 , Q_6 )$ $\bullet$ $\mathit{index} = 29,$ $( 4H - 3Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , H - Q_1 - Q_6 , H - Q_1 - Q_7 )$ $\bullet$ $\mathit{index} = 30,$ $( 8H - 3Q_1 - 3Q_2 - 3Q_3 - 3Q_4 - 3Q_5 - 3Q_6 - 3Q_7 , 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - 2Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - 2Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - 2Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - Q_4 - 2Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - 2Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - 2Q_7 )$ $\bullet$ $\mathit{index} = 31,$ $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 , Q_6 , Q_7 , Q_8 )$ $\bullet$ $\mathit{index} = 32,$ $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_5 , Q_6 , Q_8 , Q_7 )$ $\bullet$ $\mathit{index} = 33,$ $( H , Q_1 , Q_2 , Q_3 , Q_4 , Q_6 , Q_5 , Q_8 , Q_7 )$ $\bullet$ $\mathit{index} = 34,$ $( H , Q_1 , Q_2 , Q_4 , Q_3 , Q_6 , Q_5 , Q_8 , Q_7 )$ $\bullet$ $\mathit{index} = 35,$ $( H , Q_2 , Q_1 , Q_4 , Q_3 , Q_6 , Q_5 , Q_8 , Q_7 )$ $\bullet$ $\mathit{index} = 36,$ $( 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , 2H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , Q_6 , Q_7 , Q_8 )$ $\bullet$ $\mathit{index} = 37,$ $( 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , 2H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , Q_6 , Q_8 , Q_7 )$ $\bullet$ $\mathit{index} = 38,$ $( 4H - 3Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , H - Q_1 - Q_2 , H - Q_1 - Q_3 , H - Q_1 - Q_4 , H - Q_1 - Q_5 , H - Q_1 - Q_6 , H - Q_1 - Q_7 , Q_8 )$ $\bullet$ $\mathit{index} = 39,$ $( 8H - 3Q_1 - 3Q_2 - 3Q_3 - 3Q_4 - 3Q_5 - 3Q_6 - 3Q_7 , 3H - 2Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - 2Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - 2Q_3 - Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - 2Q_4 - Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - Q_4 - 2Q_5 - Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - 2Q_6 - Q_7 , 3H - Q_1 - Q_2 - Q_3 - Q_4 - Q_5 - Q_6 - 2Q_7 , Q_8 )$ $\bullet$ $\mathit{index} = 40,$ $( 17H - 6Q_1 - 6Q_2 - 6Q_3 - 6Q_4 - 6Q_5 - 6Q_6 - 6Q_7 - 6Q_8 , 6H - 3Q_1 - 2Q_2 - 2Q_3 - 2Q_4 - 2Q_5 - 2Q_6 - 2Q_7 - 2Q_8 , 6H - 2Q_1 - 3Q_2 - 2Q_3 - 2Q_4 - 2Q_5 - 2Q_6 - 2Q_7 - 2Q_8 , 6H - 2Q_1 - 2Q_2 - 3Q_3 - 2Q_4 - 2Q_5 - 2Q_6 - 2Q_7 - 2Q_8 , 6H - 2Q_1 - 2Q_2 - 2Q_3 - 3Q_4 - 2Q_5 - 2Q_6 - 2Q_7 - 2Q_8 , 6H - 2Q_1 - 2Q_2 - 2Q_3 - 2Q_4 - 3Q_5 - 2Q_6 - 2Q_7 - 2Q_8 , 6H - 2Q_1 - 2Q_2 - 2Q_3 - 2Q_4 - 2Q_5 - 3Q_6 - 2Q_7 - 2Q_8 , 6H - 2Q_1 - 2Q_2 - 2Q_3 - 2Q_4 - 2Q_5 - 2Q_6 - 3Q_7 - 2Q_8 , 6H - 2Q_1 - 2Q_2 - 2Q_3 - 2Q_4 - 2Q_5 - 2Q_6 - 2Q_7 - 3Q_8 )$

# References

Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. ISSN 0747-7171. doi: 10.1006/jsco.1996.0125. URL http://dx.doi.org/10.1006/jsco.1996.0125. Computational algebra and number theory (London, 1993).

Nicolas Bourbaki. *Lie groups and Lie algebras. Chapters 4–6.* Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. ISBN 3-540-42650-7. Translated from the 1968 French original by Andrew Pressley.

Willem A. de Graaf. *Lie algebras: theory and algorithms*, volume 56 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 2000. ISBN 0-444-50116-9.

Igor Dolgachev. *Classical Algebraic Geometry: A Modern View*. 2012. ISBN 1-107-01765-3. URL http://www.math.lsa.umich.edu/~idolga/CAG.pdf.

Patrick Du Val. On isolated singularities of surfaces which do not affect the conditions of adjunction. I,II,III. *Proc. Camb. Philos. Soc.*, 30:453–491, 1934.

Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. ISBN 0-471-32792-1. Pure and Applied Mathematics.

N. Lubbes. Phd thesis: minimal families of curves on surfaces. pages 1–274, 2011.

Ju. I. Manin. Rational surfaces over perfect fields. *Inst. Hautes Études Sci. Publ. Math.*, (30):55–113, 1966.

Yu. I. Manin. *Cubic forms*, volume 4 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, second edition, 1986. ISBN 0-444-87823-8. Algebra, geometry, arithmetic, Translated from the Russian by M. Hazewinkel.

T. Oshima. Classification of subsystems of a root system. *http://arxiv.org/abs/math/0611904*, 2006.

Robert Silhol. *Real algebraic surfaces*, volume 1392 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1989. ISBN 3-540-51563-1.

W. A. Stein et al. *Sage Mathematics Software*. The Sage Development Team, 2012. `http://www.sagemath.org`.

C. T. C. Wall. Real forms of smooth del Pezzo surfaces. *J. Reine Angew. Math.*, 375/376:47–66, 1987. ISSN 0075-4102. doi: 10.1515/crll.1987.375-376.47. URL `http://dx.doi.org/10.1515/crll.1987.375-376.47`.

**Address of author:**

King Abdullah University of Science and Technology, Thuwal, Kingdom of Saudi Arabia
**email:** niels.lubbes@gmail.com

# Index

basis, 7

C1 category, 8
C1 kit, 8
C1 label, 8
  geometric, 9
C1 label elements, 8
C1 label elements to root function, 8
C1 object, 15
C1 root system, 8
Cartan matrix, 7
category of root subsystems, 7
category of root systems, 7
complement root subsystem, 9
Cremona equivalence relation of weak Del Pezzo surfaces, 15
Cremona equivalent, 15

degree, 12
Del Pezzo one-set, 13
Del Pezzo two-set, 14
Del Pezzo zero-set, 13
double complement root subsystem, 9
Dynkin diagram, 7

enhanced Picard group, 15

geometric, 9

indecomposable class, 13
indecomposable Del Pezzo one-set, 14
indecomposable Del Pezzo zero-set, 13
irreducible Del Pezzo two-set, 14

maximal, 7

positive, 7

real structure, 22
real weak Del Pezzo surface, 22
root subsystem, 7
root system, 7
  basis, 7
  complement root subsystem, 9
  double complement root subsystem, 9
  Weyl group, 7

simple, 7
standard Del Pezzo basis, 12
standard geometrically ruled basis, 12
surface
  real structure, 22

weak Del Pezzo surface, 12
Weyl group, 7